

# Spectroscopy of $q^3\bar{q}^3$ -States in Quark Model and Baryon-Antibaryon Enhancements

Gui-Jun Ding<sup>a</sup>, Jialun Ping<sup>b</sup>, and Mu-Lin Yan<sup>a</sup>

<sup>a</sup> *Interdisciplinary Center for Theoretical Study,*

*University of Science and Technology of China, Hefei, Anhui 230026, China*

<sup>b</sup>*Department of Physics, Nanjing Normal University, Nanjing, 210097, China*

## Abstract

We study the mass spectrum of the  $q^3\bar{q}^3$  mesons both from common  $S$  wave quark model with colormagnetic interactions, and from the quark model with triquark correlations. Two cluster configurations  $(q^3) - (\bar{q}^3)$  and  $(q^2\bar{q}) - (q\bar{q}^2)$  are considered. In the spectrums of both models, we find rather stable states which have the same quantum numbers as the particle resonances which are corresponding to the  $p\bar{p}$  enhancement,  $p\bar{\Lambda}$  enhancement and  $\Lambda\bar{\Lambda}$  enhancement with spin-**0** or **1**. This implies these enhancements are NOT experimental artifacts. The color-spin-flavor structures of  $p\bar{p}$ ,  $p\bar{\Lambda}$ , and  $\Lambda\bar{\Lambda}$  enhancements are revealed. It has been found that if there exists quark correlation in the baryonium, the  $p\bar{p}$  enhancement as a baryonium is mainly a state of  $|120_{cs}, \bar{6}_c, 2, 3_f \oplus \bar{6}_f\rangle |120_{cs}, 6_c, 2, \bar{3}_f \oplus 6_f\rangle$  in the configuration  $(q^2\bar{q}) - (q\bar{q}^2)$ , with few mixture of  $|70_{cs}, 8_c, 2, 8_f\rangle |70_{cs}, 8_c, 2, 8_f\rangle$  in the configuration  $(q^3) - (\bar{q}^3)$ , where the subscripts  $cs$ ,  $c$  and  $f$  indicate colorspin, color and flavor respectively. The existence of spin-**1**  $\Lambda\bar{\Lambda}$ ,  $p\bar{\Lambda}$ ,  $p\bar{p}$  enhancements is predicted.

PACS numbers: 12.38.-t, 12.39.Jh, 12.40.Yx

## I. INTRODUCTION

There has recently been renewed theoretical interest in hadronic states composed of more exotic quark configurations than the usual  $q\bar{q}$  for mesons and  $qqq$  for baryons. The interests has been stimulated by the experimental discovery of many exotic states, e.g.  $\Theta^+$ ,  $D_s(2317)$ ,  $X(3872)$ [1]. Moreover, in the recent years, some baryon-antibaryon invariant mass threshold enhancements have been observed in experiments[2-8]. In order to interpret the  $p\bar{p}$  enhancement experiment[2], the authors of Ref.[9] pointed out that the  $q^3\bar{q}^3$  baryonium states should exist. This claim comes from analyzing the interaction among the quarks(antiquarks) in terms of the colormagnetic hyperfine interaction[18]:

$$H' = - \sum_{i>j} C_{ij} \vec{\sigma}_i \cdot \vec{\sigma}_j \vec{\lambda}_i \cdot \vec{\lambda}_j \quad (1)$$

where  $\vec{\sigma}$  and  $\vec{\lambda}$  are the Pauli matrices and Gell-Mann matrices, and the indices  $i$  and  $j$  run over all the constituent quarks and antiquarks. This analysis has been further studied by means of the chiral soliton model[11, 12]. And a remarkable prediction on the baryonium's mesonic decays from this picture in Ref.[12] has been supported by the recent BES experiment[3] significantly. In this case, because of both the experiment prompt and lack of a systemic study of the  $q^3\bar{q}^3$  system in the quark model with colormagnetic interaction in the literature, we try to provide such an investigation to the spectroscopy of the exotic ( $q^3\bar{q}^3$ ) configurations in quark model in this work.

The  $p\bar{p}$  enhancement is observed in  $J/\psi \rightarrow \gamma p\bar{p}$  [2][3], and  $\psi' \rightarrow \pi^0 p\bar{p}, \eta p\bar{p}$  [4]. The  $p\bar{\Lambda}$  enhancement has been observed in  $J/\psi \rightarrow p\bar{\Lambda}K^-$  and  $\psi' \rightarrow p\bar{\Lambda}K^-$  [5]. In B decays,  $p\bar{p}$ ,  $p\bar{\Lambda}$  and  $\Lambda\bar{\Lambda}$  enhancements also were observed, such as  $B^\pm \rightarrow p\bar{p}K^\pm$ [6],  $B^0 \rightarrow p\bar{\Lambda}\pi^-$ [7],  $B^+ \rightarrow \Lambda\bar{\Lambda}K^+$ [8]. In Ref.[2], the data of  $p\bar{p}$  invariant mass threshold enhancement has been fitted with the  $S$  and  $P$  wave Breit-Wigner resonance function, and the enhancement has naively been attributed to existence of a  $(p\bar{p})$ -baryonium resonance. Namely, the enhancement corresponds to a baryonium particle. The theoretical studies on this picture are in Refs.[9, 10, 11, 12], and an experiment recheck on it is in Ref.[3]. Noting that in order to fix the total angular momentum of the baryoniums the partial wave analysis has to be done, and the  $S$  wave or  $P$  wave data fit can not fix the spin of the baryonium. For instance, even though the baronium is fitted out by  $S$  wave, the total angular momentum of it can still be non-zero because quark's intrinsic spin is  $1/2$  and the particle is a multi-

quark system. For the sake of illustrating convenience, we will call the baryonium particle revealed by the possible enhancement effect measurements as *enhancement-baryonium*, or shortly *enhancement* hereafter. There are many interesting phenomena and puzzles in the baryon-antibaryon physics. In this paper, we mainly focus on these exotic states observed in the baryon antibaryon final states. Because the S wave fit is favored over the P wave fit, we do not consider the radial excitations in enhancements, and the models used in this paper are *S* wave type quark models, i.e., the relative angular momentum between any two quarks in the models is  $\mathbf{0}$ .

Because the baryoniums finally decay into a baryon octet state plus an antibaryon octet state and  $\mathbf{8}_f \otimes \mathbf{8}_f = \mathbf{1}_f \oplus \mathbf{8}_f \oplus \mathbf{8}_f \oplus \mathbf{10}_f \oplus \overline{\mathbf{10}}_f \oplus \mathbf{27}_f$ , these enhancements can only possibly belong to flavor  $\mathbf{1}_f$ -plet,  $\mathbf{8}_f$ -plet,  $\mathbf{10}_f$ -plet,  $\overline{\mathbf{10}}_f$ -plet or  $\mathbf{27}_f$ -plet. And the quantum numbers of the  $p\bar{p}$  enhancement most possibly are  $J^{PC} = 0^{-+}$ , so it must consist of  $q\bar{q}$ ,  $q^3\bar{q}^3$ ,  $q^5\bar{q}^5$  and so on, if the relative angular momentum between quarks equals  $\mathbf{0}$ . They are not possibly the ordinary mesonic states, then they at least are composed of  $\mathbf{3}$  quarks and  $\mathbf{3}$  antiquarks. The theoretical interpretation of these exotic state is a great challenge and many proposals have been suggested [9, 10, 11, 12, 13]. Some of them studied one particular state [9, 11, 12], others propose the possible nature of these states or the possible classification of these baryon-antibaryon enhancements[13]. In this work, we discuss the possible states of the baryon-antibaryon system (or the  $q^3\bar{q}^3$  system) and try to give a unified interpretation of these baryon-antibaryon enhancements in common quark model and in quark model with quark correlation. Since all these states have been observed in charmonium decay and B decay, it is possible that these states possess the same nature. It should be interesting to interpret them in a uniform framework.

Color plays important role in multiquark states. The configuration of color is uniquely fixed in the usual  $q\bar{q}$  mesons and baryons consisting of  $qqq$ . In the multiquark system, however, the color degree of freedom is "unfrozen" and there exist a great variety of color configurations to form color singlet states, e.g., in the system of the  $q^2\bar{q}^2$ , the two quark can be either in color  $\overline{\mathbf{3}}_c$  or  $\mathbf{6}_c$  state combining with the two antiquarks in color  $\mathbf{3}_c$  or  $\overline{\mathbf{6}}_c$  state to give a singlet[14]. Many works also have been done in exploring this point of view in pentaquark study[17]. This "color chemistry" idea is very fascinating. Even though the question that how to cluster the quarks into diquark, triquark or other structures in the systems by means of the fundamental QCD dynamics remains to be open so far, the quark-

correlation has been proved to be a very useful and important conception in the physics of multiquark state[15, 16].

We will first discuss this system in common quark model where 3 quarks and 3 antiquarks are in relative S wave and there is not correlation among the quarks(antiquarks). Under the QCD requirement that the baryoniums have to be in color singlets, the spectrum of enhancement has been calculated, and the eigenstates are linear combinations of states which are in different  $SU(6)_{cs}$  representations. For the sake of completeness, we then discuss the effect of quark correlation in the baryinoim. There are many different ways to form color singlet for the  $q^3\bar{q}^3$  system, and we expect the mesonic states corresponding to the different color couplings between the quark subsystems have different physical properties. We consider two case in this work, the first case is the  $(q^3) - (\bar{q}^3)$  configuration and another configuration is  $(q^2\bar{q}) - (q\bar{q}^2)$ . We assume the mass splitting is due to the colormagnetic interaction [18], we also list the possible states and the mass splitting in the two cases. Since the S wave fit is favored, we assume the relative angular momentum of the three quarks(antiquarks) in each cluster is zero, and the two clusters are also in relative S wave.

In this work we take a close look at the  $q^3\bar{q}^3$  system in quark model with the mass splitting due to colormagnetic hyperfine interaction, and we don't consider the effect of  $SU(3)_f$  flavor symmetry breaking, and the results are qualitatively correct.

The paper is organized as followings: in the section II, we review the description of colormagnetic hyperfine interaction between quarks in hadrons and its application to the  $q^2\bar{q}^2$  system; Section III is devoted to study the  $q^3\bar{q}^3$  system in common quark model; In Section IV, we study two interesting configurations with quark correlation:  $(q^3) - (\bar{q}^3)$  and  $(q^2\bar{q}) - (\bar{q}^2q)$ ; Finally, we briefly summary the results and give some discussions. In Appendix A, we show an example to deal with the mixing of different  $SU(6)_{cs}$  multiplets under colormagnetic interaction, and we also list the matrix form of the hamiltonian  $H'$  in the subspace expanded by the states with definite spin in each case in Appendix B.

## II. COLORMAGNETIC HYPERFINE INTERACTION AND ITS APPLICATION TO THE $q^2\bar{q}^2$ MESONS

The spectroscopically important interaction between quarks(aside from the interaction which provides the confinement and sets the overall scale) is the spin-spin force mediated

by one gluon exchange. The interaction hamiltonian  $H'$  between quarks is given by Eq(1). Generally the coefficients  $C_{ij}$  are dependent on the quark masses of flavor  $i$  and  $j$  and the properties of the spatial wave functions of the quarks and the antiquarks in the system and parameterize the strength of the interaction. Following the common colormagnetic hyperfine studies in literature [17, 19], we take  $C_{ij} = C$  neglecting the breaking of the flavor  $SU(3)_f$  symmetry. It is straightforward to express the colormagnetic interaction in terms of the quadratic Casimir operators of  $SU(2)_s$ ,  $SU(3)_c$  and  $SU(6)_{cs}$  of the total system, the quarks and the antiquarks,

$$H' = C[8N + 2C_6(tot) - \frac{4}{3}S_{tot}(S_{tot} + 1) - 2C_3(tot) + 4C_3(Q) + \frac{8}{3}S_Q(S_Q + 1) - 4C_6(Q) + 4C_3(\bar{Q}) + \frac{8}{3}S_{\bar{Q}}(S_{\bar{Q}} + 1) - 4C_6(\bar{Q})] \quad (2)$$

Where  $C_3$  and  $C_6$  are the quadratic Casimir operator of the  $SU(3)_c$  and  $SU(6)_{cs}$ ,  $S$  is the spin, the labels  $Q$ ,  $\bar{Q}$  and  $tot$  refer to the quarks, antiquarks, and the entire system respectively, and  $N$  is the total number of the quarks and antiquarks in the system, for the  $q^3\bar{q}^3$  system  $N = 6$ . The Casimir operators are normalized conventionally, e.g.  $C_3(3) = \frac{4}{3}$ ,  $C_6(6) = \frac{35}{6}$ . Eigenstates of total  $SU(6)_{cs}$  for  $q^3\bar{q}^3$  are generally mixtures of the states with different color and spin representations of quarks and antiquarks respectively, and the colormagnetic interaction usually mixes states of different total colorspin.

So far various systems has been discussed in quark model with colormagnetic interaction. It was first considered in Ref.[18], where it was applied to study the mass splittings of the S wave color singlet  $q\bar{q}$  mesons and  $q^3$  baryons, and it gives a good description of the spectrums. And Jaffe studied the  $q^2\bar{q}^2$  system and the  $0^+$  mesonic states with mass bellow 1GeV, i.e.,  $a_0(980)$ ,  $f_0(980)$  are identified as four quarks states [19]. Dibaryon( the  $q^6$  system ) also was discussed [20]. Recently pentaquark (the  $q^4\bar{q}$  system) has been widely studied in colormagnetic interaction models [17].

In order to clarify the notations and methods in the studies of colomagnetic interaction, we briefly review Jaffe's works on  $q^2\bar{q}^2$ [19, 22]. Because of the Fermi statistics, the allowed states of the 2 quarks are

$$|\mathbf{21}_{cs}, \mathbf{\bar{3}}_f\rangle, \quad |\mathbf{15}_{cs}, \mathbf{6}_f\rangle \quad (3)$$

The 2 antiquarks are in conjugate representations,

$$|\mathbf{\bar{21}}_{cs}, \mathbf{3}_f\rangle, \quad |\mathbf{\bar{15}}_{cs}, \mathbf{\bar{6}}_f\rangle \quad (4)$$

where the first number in the bracket is the dimension of the irreducible representation of  $SU(6)_{cs}$  and the second number is the dimension of the irreducible representation of  $SU(3)_f$ . The physical  $q^2\bar{q}^2$  states are color singlets, and there are four combinations:

$$\begin{aligned}
\mathbf{21}_{cs} \otimes \overline{\mathbf{21}}_{cs} &= \mathbf{1}_{cs} \oplus \mathbf{35}_{cs} \oplus \mathbf{405}_{cs}, \quad \text{with flavor in } \overline{\mathbf{3}}_f \otimes \mathbf{3}_f = \mathbf{1}_f \oplus \mathbf{8}_f \\
\mathbf{15}_{cs} \otimes \overline{\mathbf{15}}_{cs} &= \mathbf{1}_{cs} \oplus \mathbf{35}_{cs} \oplus \mathbf{189}_{cs}, \quad \text{with flavor in } \mathbf{6}_f \otimes \overline{\mathbf{6}}_f = \mathbf{1}_f \oplus \mathbf{8}_f \oplus \mathbf{27}_f \\
\mathbf{21}_{cs} \otimes \overline{\mathbf{15}}_{cs} &= \mathbf{35}_{cs} \oplus \mathbf{280}_{cs}, \quad \text{with flavor in } \overline{\mathbf{3}}_f \otimes \overline{\mathbf{6}}_f = \mathbf{8}_f \oplus \overline{\mathbf{10}}_f \\
\mathbf{15}_{cs} \otimes \overline{\mathbf{21}}_{cs} &= \overline{\mathbf{35}}_{cs} \oplus \overline{\mathbf{280}}_{cs}, \quad \text{with flavor in } \mathbf{6}_f \otimes \mathbf{3}_f = \mathbf{8}_f \oplus \mathbf{10}_f
\end{aligned} \tag{5}$$

The four sector's multiplets are not mixed by the colormagnetic interaction.

For the  $\mathbf{21}_{cs} \otimes \overline{\mathbf{21}}_{cs}$  case, the  $SU(6)_{cs}$  decomposition with respect to  $SU(3)_c \otimes SU(2)_s$  reveals the following color singlets:

$$\begin{aligned}
(\mathbf{1}_c, \mathbf{1}) &\subset \mathbf{1}_{cs} \\
(\mathbf{1}_c, \mathbf{3}) &\subset \mathbf{35}_{cs} \\
(\mathbf{1}_c, \mathbf{1}) \text{ and } (\mathbf{1}_c, \mathbf{5}) &\subset \mathbf{405}_{cs}
\end{aligned} \tag{6}$$

The wavefunctions of the spin-2 states  $|\mathbf{405}_{cs}, \mathbf{1}_c, \mathbf{5}; \mathbf{3}_f \otimes \overline{\mathbf{3}}_f\rangle$  and spin-1  $|\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{3}_f \otimes \overline{\mathbf{3}}_f\rangle$  respectively are,

$$\begin{aligned}
|\mathbf{405}_{cs}, \mathbf{1}_c, \mathbf{5}; \mathbf{3}_f \otimes \overline{\mathbf{3}}_f\rangle &= |(\mathbf{21}_{cs}, \mathbf{6}_c, \mathbf{3}; \overline{\mathbf{3}}_f), (\overline{\mathbf{21}}_{cs}, \overline{\mathbf{6}}_c, \mathbf{3}; \mathbf{3}_f), \mathbf{1}_c, \mathbf{5}; \mathbf{3}_f \otimes \overline{\mathbf{3}}_f\rangle \\
|\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{3}_f \otimes \overline{\mathbf{3}}_f\rangle &= |(\mathbf{21}_{cs}, \mathbf{6}_c, \mathbf{3}; \overline{\mathbf{3}}_f), (\overline{\mathbf{21}}_{cs}, \overline{\mathbf{6}}_c, \mathbf{3}; \mathbf{3}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{3}_f \otimes \overline{\mathbf{3}}_f\rangle
\end{aligned} \tag{7}$$

The corresponding colormagnetic interaction energy are  $\frac{32}{3}$  and  $-16$  respectively. Here the wave function is expressed in terms of the conventional notations[27]:

$$\begin{aligned}
&|D_{cs}(tot), D_c(tot), 2S(tot) + 1; D_f(tot)\rangle \\
&= |(D_{cs}(Q), D_c(Q), 2S(Q) + 1; D_f(Q)), (D_{cs}(\overline{Q}), D_c(\overline{Q}), 2S(\overline{Q}) + 1; D_f(\overline{Q})), \\
&D_c(tot), 2S(tot) + 1; D_f(tot)\rangle,
\end{aligned}$$

where  $D_{cs}(tot), D_{cs}(Q), D_{cs}(\overline{Q})$  denote the dimension of the irreducible  $SU(6)_{cs}$  representation of the total system, the quarks, and the antiquarks respectively. And similarly for the notations  $D_c(id), D_f(id)$ , the  $id$  represents  $tot, Q$  or  $\overline{Q}$ .  $S(tot), S(Q), S(\overline{Q})$  are respectively the spin of the total system, the quarks and the antiquarks.

The two spin-0 states are linear combinations of  $(\bar{\mathbf{3}}_c, \mathbf{1}) \otimes (\mathbf{3}_c, \mathbf{1})$  with  $(\mathbf{6}_c, \mathbf{3}) \otimes (\bar{\mathbf{6}}_c, \mathbf{3})$ ,

$$\begin{aligned}
|1_{cs}, 1_c, \mathbf{1}; \mathbf{3}_f \otimes \bar{\mathbf{3}}_f\rangle &= \sqrt{\frac{6}{7}} |(\mathbf{21}_{cs}, \mathbf{6}_c, \mathbf{3}; \bar{\mathbf{3}}_f), (\bar{\mathbf{21}}_{cs}, \bar{\mathbf{6}}_c, \mathbf{3}; \mathbf{3}_f), 1_c, \mathbf{1}; \mathbf{3}_f \otimes \bar{\mathbf{3}}_f\rangle \\
&+ \sqrt{\frac{1}{7}} |(\mathbf{21}_{cs}, \bar{\mathbf{3}}_c, \mathbf{1}; \bar{\mathbf{3}}_f), (\bar{\mathbf{21}}_{cs}, \mathbf{3}_c, \mathbf{1}; \mathbf{3}_f), 1_c, \mathbf{1}; \mathbf{3}_f \otimes \bar{\mathbf{3}}_f\rangle \\
|405_{cs}, 1_c, \mathbf{1}; \mathbf{3}_f \otimes \bar{\mathbf{3}}_f\rangle &= \sqrt{\frac{1}{7}} |(\mathbf{21}_{cs}, \mathbf{6}_c, \mathbf{3}; \bar{\mathbf{3}}_f), (\bar{\mathbf{21}}_{cs}, \bar{\mathbf{6}}_c, \mathbf{3}; \mathbf{3}_f), 1_c, \mathbf{1}; \mathbf{3}_f \otimes \bar{\mathbf{3}}_f\rangle \\
&- \sqrt{\frac{6}{7}} |(\mathbf{21}_{cs}, \bar{\mathbf{3}}_c, \mathbf{1}; \bar{\mathbf{3}}_f), (\bar{\mathbf{21}}_{cs}, \mathbf{3}_c, \mathbf{1}; \mathbf{3}_f), 1_c, \mathbf{1}; \mathbf{3}_f \otimes \bar{\mathbf{3}}_f\rangle
\end{aligned} \tag{8}$$

The colormagnetic interaction mixes these two states, and the eigenstates of  $H'$  (Eq.(2)) are

$$\begin{aligned}
|\mathbf{1}, \mathbf{3}_f \otimes \bar{\mathbf{3}}_f\rangle_1 &= 0.972 |1_{cs}, 1_c, \mathbf{1}; \mathbf{3}_f \otimes \bar{\mathbf{3}}_f\rangle + 0.233 |405_{cs}, 1_c, \mathbf{1}; \mathbf{3}_f \otimes \bar{\mathbf{3}}_f\rangle \\
|\mathbf{1}, \mathbf{3}_f \otimes \bar{\mathbf{3}}_f\rangle_2 &= 0.233 |1_{cs}, 1_c, \mathbf{1}; \mathbf{3}_f \otimes \bar{\mathbf{3}}_f\rangle - 0.972 |405_{cs}, 1_c, \mathbf{1}; \mathbf{3}_f \otimes \bar{\mathbf{3}}_f\rangle
\end{aligned} \tag{9}$$

The colormagnetic interaction energies ( i.e. the eigenvalues of  $H'$  ) are -43.26 and -19.37 respectively. The color singlet states and the corresponding colormagnetic energies for the other three sectors in Eq.(5) have also been derived in the same way by Jaffe[19]. Thus, the color singlet spectroscopy of  $q^2\bar{q}^2$  states in quark model has been revealed. Among all of these multiplets, the nonet multiplet  $|\mathbf{1}, \mathbf{3}_f \otimes \bar{\mathbf{3}}_f\rangle_1$  has the largest colormagnetic binding energy. Therefore, they are the lightest states, and one can physically regard them as the known  $0^+$  nonet mesons below 1 GeV. All of these have been done and be well known. An interesting question and task is how to extent Jaffe's analysis on  $q^2\bar{q}^2$ -system in quark model into the  $q^3\bar{q}^3$ -system, and to explore the corresponding physics. In the following of this paper, we perform such an investigation.

### III. THE $q^3\bar{q}^3$ SYSTEM AND THE ENHANCEMENT FROM THE COMMON QUARK MODEL

Since the experimental discovery of the enhancements which can be regarded as baryoniums, it is necessary to explore the  $q^3\bar{q}^3$  system in the framework of colormagnetic interaction. Now let us to extent the Jaffe's studies on  $q^2\bar{q}^2$  system to  $q^3\bar{q}^3$  one. Due to the increase of a pair of quark and antiquark, the dimension of the irreducible representation will become rather large, and more techniques and skills of group theory are needed for dealing with this multi-quark system. The calculations are rather lengthy. In order to ensure the correctness of calculation results, we have checked each analytical group theory derivations by the

computer program calculations. The results obtained by the two approaches are exactly the same.

Similar to studying the usual  $q\bar{q}$  meson and  $q^3$  baryon from the common quark model, we study the  $q^3\bar{q}^3$  system in the quark model in order to understand the observed baryon antibaryon enhancements, and all quarks are in relative S wave. The configuration is schematically shown in Fig.3.

Because of the Fermi statistics, the three quarks's state must be antisymmetric in the combined wavefunctions of flavor $\otimes$ color $\otimes$ spin $\otimes$ space. Because the quarks are in a relative S wave, the flavor $\otimes$ color $\otimes$ spin wavefunction must be antisymmetric. Under the group  $SU(18) \supset SU(6)_{cs} \otimes SU(3)_f \supset SU(3)_c \otimes SU(2)_s \otimes SU(3)_f$ , the three quarks must be in a completely antisymmetric state and it therefore transform under the  $\frac{18 \cdot 17 \cdot 16}{3!} = \mathbf{816}$  dimensional irreducible representation of  $SU(18)$ . The reduction of the  $SU(18)$  irreducible representation  $\mathbf{816}$  with respect to  $SU(6)_{cs} \otimes SU(3)_f$  is [24]

$$\mathbf{816} = (\mathbf{70}_{cs}, \mathbf{8}_f) + (\mathbf{56}_{cs}, \mathbf{1}_f) + (\mathbf{20}_{cs}, \mathbf{10}_f), \quad (10)$$

Here the notation is the same as Eq.(3), and the allowed states of the 3 quarks are

$$|\mathbf{70}_{cs}, \mathbf{8}_f\rangle, \quad |\mathbf{56}_{cs}, \mathbf{1}_f\rangle, \quad |\mathbf{20}_{cs}, \mathbf{10}_f\rangle, \quad (11)$$

And the 3 antiquarks are in conjugate representations,

$$|\overline{\mathbf{70}}_{cs}, \mathbf{8}_f\rangle, \quad |\overline{\mathbf{56}}_{cs}, \mathbf{1}_f\rangle, \quad |\overline{\mathbf{20}}_{cs}, \overline{\mathbf{10}}_f\rangle. \quad (12)$$

Next we combine the states of the 3 quarks in Eq.(11) and that of the 3 antiquarks in Eq.(12) to obtain the states of the total systems which is color singlet, in the following we consider the allowed states of the total system and the corresponding colormagnetic energy case by case. In the text, we mainly list the results and illustrate calculation skills, and in Appendix A, we take the mixing of the three spin-1 states in the case of  $\mathbf{70}_{cs} \otimes \mathbf{20}_{cs}$  as an example in order to show how analytical calculations are performed in details. In Appendix B, we write the explicit matrix form of the interaction hamiltonian  $H'$  in the subspace expanded by the states of definite spin in each case, from which we can obtain the eigenstates and the corresponding colormagnetic energies.

#### 1. $\mathbf{20}_{cs} \otimes \overline{\mathbf{20}}_{cs}$



From Eq.(11) and Eq.(12), the multiplets must be in the flavor representation  $\mathbf{10}_f \otimes \overline{\mathbf{10}}_f = \mathbf{64}_f \oplus \mathbf{27}_f \oplus \mathbf{8}_f \oplus \mathbf{1}_f$ , from Fig.1(VI) we can see

$$\mathbf{20}_{cs} \otimes \overline{\mathbf{20}}_{cs} = \mathbf{175}_{cs} \oplus \mathbf{189}_{cs} \oplus \mathbf{35}_{cs} \oplus \mathbf{1}_{cs} \quad (13)$$

The  $\text{SU}(6)_{cs}$  decomposition with respect to  $\text{SU}(3)_c \otimes \text{SU}(2)_s$  which are listed in Table VII and Table VIII shows the following color singlet,

$$\begin{aligned} (\mathbf{1}_c, \mathbf{1}) &\subset \mathbf{1}_{cs} \\ (\mathbf{1}_c, \mathbf{3}) &\subset \mathbf{35}_{cs} \\ (\mathbf{1}_c, \mathbf{3}) \text{ and } (\mathbf{1}_c, \mathbf{7}) &\subset \mathbf{175}_{cs} \\ (\mathbf{1}_c, \mathbf{1}) \text{ and } (\mathbf{1}_c, \mathbf{5}) &\subset \mathbf{189}_{cs} \end{aligned} \quad (14)$$

In order to derive the mass splitting due to  $H'$  by using Eq.(2), we must know which representations of the  $\text{SU}(3)_c$  and of  $\text{SU}(2)_s$  the quarks and antiquarks in the system belong to. In the followings, we study the  $H'$  eigenstates with spin-**3**, -**1** and -**0** respectively, and their corresponding eigenvalues:

**A)**, The spin-**3** states  $|\mathbf{175}_{cs}, \mathbf{1}_c, \mathbf{7}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle$  and the spin-**2** states  $|\mathbf{189}_{cs}, \mathbf{1}_c, \mathbf{5}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle$  are trivial. The wavefunction of the former is

$$|\mathbf{175}_{cs}, \mathbf{1}_c, \mathbf{7}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle = |(\mathbf{20}_{cs}, \mathbf{1}_c, \mathbf{4}; \mathbf{10}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{1}_c, \mathbf{4}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{7}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle \quad (15)$$

Since

$$\begin{aligned} C_6(tot) &= C_6(\mathbf{175}_{cs}) = 24, \quad S_{tot} = \mathbf{3}, \quad C_3(tot) = C_3(\mathbf{1}_c) = 0, \\ C_3(Q) &= C_3(\mathbf{1}_c) = 0, \quad S_Q = \frac{3}{2}, \quad C_6(Q) = C_6(\mathbf{20}_{cs}) = \frac{21}{2}, \\ C_3(\overline{Q}) &= C_3(\mathbf{1}_c) = 0, \quad S_{\overline{Q}} = \frac{3}{2}, \quad C_6(\overline{Q}) = C_6(\overline{\mathbf{20}}_{cs}) = \frac{21}{2}. \end{aligned}$$

Substituting the above quantities into Eq.(2), then we get the corresponding color-magnetic energy  $16C$ . And the latter is

$$|\mathbf{189}_{cs}, \mathbf{1}_c, \mathbf{5}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle = |(\mathbf{20}_{cs}, \mathbf{1}_c, \mathbf{4}; \mathbf{10}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{1}_c, \mathbf{4}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{5}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle \quad (16)$$

The corresponding  $\text{SU}(6)$  Casimir operator is  $C_6(tot) = C_6(\mathbf{189}_{cs}) = 20$ , and the colormagnetic energy is also  $16C$ .

B), The two spin-1 states  $|\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle$  and  $|\mathbf{175}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle$  are linear combinations of the states  $|(\mathbf{20}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{10}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle$  and  $|(\mathbf{20}_{cs}, \mathbf{1}_c, \mathbf{4}; \mathbf{10}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{1}_c, \mathbf{4}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle$ . Following the method of the Appendix A, we can obtain

$$\begin{aligned}
|\mathbf{175}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle &= \sqrt{\frac{5}{9}}|(\mathbf{20}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{10}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle \\
&- \sqrt{\frac{4}{9}}|(\mathbf{20}_{cs}, \mathbf{1}_c, \mathbf{4}; \mathbf{10}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{1}_c, \mathbf{4}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle, \\
|\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle &= -\sqrt{\frac{4}{9}}|(\mathbf{20}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{10}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle \\
&- \sqrt{\frac{5}{9}}|(\mathbf{20}_{cs}, \mathbf{1}_c, \mathbf{4}; \mathbf{10}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{1}_c, \mathbf{4}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle
\end{aligned} \tag{17}$$

These two states are mixed by the colormagnetic interaction hamiltonian  $H'$ ,

$$\begin{aligned}
H'|\mathbf{175}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle &= \frac{16C}{9}(19|\mathbf{175}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle - \sqrt{5}|\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle) \\
H'|\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle &= \frac{16C}{9}(-\sqrt{5}|\mathbf{175}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle + 5|\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle)
\end{aligned}$$

From above, we can learn that the hamiltonian  $H'$  can be written explicitly as the following matrix form in subspace expanded by the two states,

$$\frac{16C}{9} \begin{pmatrix} 19 & -\sqrt{5} \\ -\sqrt{5} & 5 \end{pmatrix} \tag{18}$$

The eigenstates are

$$\begin{aligned}
|\mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle_1 &= 0.988|\mathbf{175}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle - 0.154|\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle \\
&= 0.839|(\mathbf{20}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{10}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle - 0.544|(\mathbf{20}_{cs}, \mathbf{1}_c, \mathbf{4}; \mathbf{10}_f), \\
&(\overline{\mathbf{20}}_{cs}, \mathbf{1}_c, \mathbf{4}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle,
\end{aligned} \tag{19}$$

$$\begin{aligned}
|\mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle_2 &= 0.154|\mathbf{175}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle + 0.988|\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle \\
&= -0.544|(\mathbf{20}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{10}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle - 0.839|(\mathbf{20}_{cs}, \mathbf{1}_c, \mathbf{4}; \mathbf{10}_f), \\
&(\overline{\mathbf{20}}_{cs}, \mathbf{1}_c, \mathbf{4}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle
\end{aligned} \tag{20}$$

The corresponding colormagnetic energy are 34.397C and 8.269C.

C), In the same way, we can calculate the mixing of two spin-0 states  $|\mathbf{189}_{cs}, \mathbf{1}_c, \mathbf{1}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle$  and  $|\mathbf{1}_{cs}, \mathbf{1}_c, \mathbf{1}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle$ . The eigenstates are

$$\begin{aligned} |\mathbf{1}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle_1 &= 0.996|\mathbf{189}_{cs}, \mathbf{1}_c, \mathbf{1}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle - 0.090|\mathbf{1}_{cs}, \mathbf{1}_c, \mathbf{1}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle \\ &= 0.526|(\mathbf{20}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{10}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle - 0.851|(\mathbf{20}_{cs}, \mathbf{1}_c, \mathbf{4}; \mathbf{10}_f), \\ &\quad (\overline{\mathbf{20}}_{cs}, \mathbf{1}_c, \mathbf{4}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle, \end{aligned} \quad (21)$$

$$\begin{aligned} |\mathbf{1}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle_2 &= 0.090|\mathbf{189}_{cs}, \mathbf{1}_c, \mathbf{1}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle + 0.996|\mathbf{1}_{cs}, \mathbf{1}_c, \mathbf{1}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle \\ &= -0.851|(\mathbf{20}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{10}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle - 0.526|(\mathbf{20}_{cs}, \mathbf{1}_c, \mathbf{4}; \mathbf{10}_f), \\ &\quad (\overline{\mathbf{20}}_{cs}, \mathbf{1}_c, \mathbf{4}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{10}_f \otimes \overline{\mathbf{10}}_f\rangle \end{aligned} \quad (22)$$

The colormagnetic energy separately are 25.889C and -9.889C.

## 2. $\mathbf{56}_{cs} \otimes \overline{\mathbf{20}}_{cs}$ and $\mathbf{20}_{cs} \otimes \overline{\mathbf{56}}_{cs}$

The two multiplets are related by charge conjugation, and we discuss the case of  $\mathbf{56}_{cs} \otimes \overline{\mathbf{20}}_{cs}$ , the flavor representation is  $\mathbf{1}_f \otimes \overline{\mathbf{10}}_f = \overline{\mathbf{10}}_f$ , and

$$\mathbf{56}_{cs} \otimes \overline{\mathbf{20}}_{cs} = \mathbf{840}_{cs} \oplus \mathbf{280}_{cs} \quad (23)$$

The color singlets are as followings

$$\begin{aligned} (\mathbf{1}_c, \mathbf{3}) &\subset \mathbf{280}_{cs} \\ (\mathbf{1}_c, \mathbf{1}) &\subset \mathbf{840}_{cs} \end{aligned} \quad (24)$$

The two wavefunctions  $|\mathbf{840}_{cs}, \mathbf{1}_c, \mathbf{1}; \mathbf{1}_f \otimes \overline{\mathbf{10}}_f\rangle$  and  $|\mathbf{280}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{1}_f \otimes \overline{\mathbf{10}}_f\rangle$  are trivial, and they are as follows

$$|\mathbf{840}_{cs}, \mathbf{1}_c, \mathbf{1}; \mathbf{1}_f \otimes \overline{\mathbf{10}}_f\rangle = |(\mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{1}_f \otimes \overline{\mathbf{10}}_f\rangle \quad (25)$$

$$|\mathbf{280}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{1}_f \otimes \overline{\mathbf{10}}_f\rangle = |(\mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{1}_f \otimes \overline{\mathbf{10}}_f\rangle \quad (26)$$

The colormagnetic energy respectively are 16C and  $-\frac{32}{3}$ C. In the same way we can obtain the wavefunctions in the case of  $\mathbf{20}_{cs} \otimes \overline{\mathbf{56}}_{cs}$  by interchange quarks and antiquarks in Eq.(25) and Eq.(26), and the corresponding colormagnetic energy is invariant.

## 3. $\mathbf{56}_{cs} \otimes \overline{\mathbf{56}}_{cs}$

The multiplets belong to the flavor representation  $\mathbf{1}_f \otimes \mathbf{1}_f = \mathbf{1}_f$ , and

$$\mathbf{56}_{cs} \otimes \overline{\mathbf{56}}_{cs} = \mathbf{2695}_{cs} \oplus \mathbf{405}_{cs} \oplus \mathbf{35}_{cs} \oplus \mathbf{1}_{cs} \quad (27)$$

The overall color singlets are as followings,

$$\begin{aligned} (\mathbf{1}_c, \mathbf{1}) &\subset \mathbf{1}_{cs} \\ (\mathbf{1}_c, \mathbf{3}) &\subset \mathbf{35}_{cs} \\ (\mathbf{1}_c, \mathbf{1}) \text{ and } (\mathbf{1}_c, \mathbf{5}) &\subset \mathbf{405}_{cs} \\ (\mathbf{1}_c, \mathbf{3}) \text{ and } (\mathbf{1}_c, \mathbf{7}) &\subset \mathbf{2695}_{cs} \end{aligned} \quad (28)$$

A), The spin-2 and spin-3 states are trivial, their wavefunctions separately are

$$|\mathbf{2695}_{cs}, \mathbf{1}_c, \mathbf{7}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle = |(\mathbf{56}_{cs}, \mathbf{10}_c, \mathbf{4}; \mathbf{1}_f), (\overline{\mathbf{56}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{4}; \mathbf{1}_f), \mathbf{1}_c, \mathbf{7}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle \quad (29)$$

$$|\mathbf{405}_{cs}, \mathbf{1}_c, \mathbf{5}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle = |(\mathbf{56}_{cs}, \mathbf{10}_c, \mathbf{4}; \mathbf{1}_f), (\overline{\mathbf{56}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{4}; \mathbf{1}_f), \mathbf{1}_c, \mathbf{5}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle \quad (30)$$

The corresponding colormagnetic energy separately are 16C and -16C.

B), The two spin-0 states  $|\mathbf{1}_{cs}, \mathbf{1}_c, \mathbf{1}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle$  and  $|\mathbf{405}_{cs}, \mathbf{1}_c, \mathbf{1}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle$  are linear combinations of the states  $|(\mathbf{56}_{cs}, \mathbf{10}_c, \mathbf{4}; \mathbf{1}_f), (\overline{\mathbf{56}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{4}; \mathbf{1}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle$  and  $|(\mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f), (\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle$ . By using the latter two states as base vectors, the matrix representation of  $H'$  can be derived explicitly (please see Appendix B, Eq.(B1)). Thus,  $H'$ 's eigenstates read

$$\begin{aligned} |1, 1_f \otimes 1_f\rangle_1 &= 0.591 |(\mathbf{56}_{cs}, \mathbf{10}_c, \mathbf{4}; \mathbf{1}_f), (\overline{\mathbf{56}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{4}; \mathbf{1}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle \\ &+ 0.807 |(\mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f), (\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle, \end{aligned} \quad (31)$$

$$\begin{aligned} |1, 1_f \otimes 1_f\rangle_2 &= 0.807 |(\mathbf{56}_{cs}, \mathbf{10}_c, \mathbf{4}; \mathbf{1}_f), (\overline{\mathbf{56}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{4}; \mathbf{1}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle \\ &- 0.591 |(\mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f), (\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle \end{aligned} \quad (32)$$

The corresponding colormagnetic energy are -82.533C and -29.467C.

C), The two spin-1 states  $|\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle$  and  $|\mathbf{2695}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle$  (or  $|(\mathbf{56}_{cs}, \mathbf{10}_c, \mathbf{4}; \mathbf{1}_f), (\overline{\mathbf{56}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{4}; \mathbf{1}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle$ ,  $|(\mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f), (\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle$ ) are mixed by the hamiltonian

$H'$ . The matrix representation of  $H'$  in the space expanded by the latter two states vector have been given by Eq.(B2) in Appendix B, then the eigenstates are

$$\begin{aligned} |\mathbf{3}, \mathbf{1}_f \otimes \mathbf{1}_f\rangle_1 &= 0.864|(\mathbf{56}_{cs}, \mathbf{10}_c, \mathbf{4}; \mathbf{1}_f), (\overline{\mathbf{56}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{4}; \mathbf{1}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle \\ &+ 0.504|(\mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f), (\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle, \end{aligned} \quad (33)$$

$$\begin{aligned} |\mathbf{3}, \mathbf{1}_f \otimes \mathbf{1}_f\rangle_2 &= -0.504|(\mathbf{56}_{cs}, \mathbf{10}_c, \mathbf{4}; \mathbf{1}_f), (\overline{\mathbf{56}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{4}; \mathbf{1}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle \\ &+ 0.864|(\mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f), (\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{1}_f \otimes \mathbf{1}_f\rangle \end{aligned} \quad (34)$$

the corresponding eigenvalue are -48.331C, -5.003C.

4.  $\mathbf{70}_{cs} \otimes \overline{\mathbf{20}}_{cs}$  and  $\mathbf{20}_{cs} \otimes \overline{\mathbf{70}}_{cs}$ .

Similar to the case 2., the multiplets are related together by charge conjugation, then we discuss  $\mathbf{70}_{cs} \otimes \overline{\mathbf{20}}_{cs}$ , and we can easily obtain the results for  $\mathbf{20}_{cs} \otimes \overline{\mathbf{70}}_{cs}$ . The states can belong to the flavor representation  $\mathbf{8}_f \otimes \overline{\mathbf{10}}_f = \overline{\mathbf{35}}_f \oplus \mathbf{27}_f \oplus \overline{\mathbf{10}}_f \oplus \mathbf{8}_f$ , and

$$\mathbf{70}_{cs} \oplus \overline{\mathbf{20}}_{cs} = \mathbf{896}_{cs} \oplus \mathbf{280}_{cs} \oplus \mathbf{189}_{cs} \oplus \mathbf{35}_{cs} \quad (35)$$

The overall color singlets are

$$\begin{aligned} (\mathbf{1}_c, \mathbf{3}) &\subset \mathbf{35}_{cs} \\ (\mathbf{1}_c, \mathbf{1}) \text{ and } (\mathbf{1}_c, \mathbf{5}) &\subset \mathbf{189}_{cs} \\ (\mathbf{1}_c, \mathbf{3}) &\subset \mathbf{280}_{cs} \\ (\mathbf{1}_c, \mathbf{3}) \text{ and } (\mathbf{1}_c, \mathbf{5}) &\subset \mathbf{896}_{cs} \end{aligned} \quad (36)$$

A), The spin-0 states are trivial, its wavefunctions are

$$|\mathbf{189}_{cs}, \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle = |(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle \quad (37)$$

The colormagnetic energy is 8C.

B), The three spin-1 states  $|\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle$ ,  $|\mathbf{280}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle$  and  $|\mathbf{896}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle$  are linear combinations of the states  $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle$ ,  $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle$  and

$|(\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), (\mathbf{20}_{cs}, \mathbf{1}_c, \mathbf{4}; \mathbf{10}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \mathbf{10}_f\rangle$  (please see Appendix A for details),

$$\begin{aligned}
|896_{cs}, 1_c, 3; 8_f \otimes \overline{10}_f\rangle &= -\sqrt{\frac{1}{9}}|(\mathbf{70}_{cs}, 8_c, 4; 8_f), (\overline{\mathbf{20}}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle \\
&- \sqrt{\frac{4}{9}}|(\mathbf{70}_{cs}, 8_c, 2; 8_f), (\overline{\mathbf{20}}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle + \sqrt{\frac{4}{9}}|(\mathbf{70}_{cs}, 1_c, 2; 8_f), \\
&(\overline{\mathbf{20}}_{cs}, 1_c, 4; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle; \\
|280_{cs}, 1_c, 3; 8_f \otimes \overline{10}_f\rangle &= -\sqrt{\frac{4}{9}}|(\mathbf{70}_{cs}, 8_c, 4; 8_f), (\overline{\mathbf{20}}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle \\
&- \sqrt{\frac{1}{9}}|(\mathbf{70}_{cs}, 8_c, 2; 8_f), (\overline{\mathbf{20}}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle - \sqrt{\frac{4}{9}}|(\mathbf{70}_{cs}, 1_c, 2; 8_f), \\
&(\overline{\mathbf{20}}_{cs}, 1_c, 4; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle; \\
|35_{cs}, 1_c, 3; 8_f \otimes \overline{10}_f\rangle &= -\sqrt{\frac{4}{9}}|(\mathbf{70}_{cs}, 8_c, 4; 8_f), (\overline{\mathbf{20}}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle \\
&+ \sqrt{\frac{4}{9}}|(\mathbf{70}_{cs}, 8_c, 2; 8_f), (\overline{\mathbf{20}}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle + \sqrt{\frac{1}{9}}|(\mathbf{70}_{cs}, 1_c, 2; 8_f), \\
&(\overline{\mathbf{20}}_{cs}, 1_c, 4; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle
\end{aligned} \tag{38}$$

These states are mixed by colormagnetic interaction  $H'$ , from above we can calculate the eigenstates

$$\begin{aligned}
|3, 8_f \otimes \overline{10}_f\rangle_1 &= 0.841|896_{cs}, 1_c, 3; 8_f \otimes \overline{10}_f\rangle + 0.536|280_{cs}, 1_c, 3; 8_f \otimes \overline{10}_f\rangle \\
&+ 0.069|35_{cs}, 1_c, 3; 8_f \otimes \overline{10}_f\rangle = -0.684|(\mathbf{70}_{cs}, 8_c, 4; 8_f), (\overline{20}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle \\
&- 0.693|(\mathbf{70}_{cs}, 8_c, 2; 8_f), (\overline{20}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle + 0.226|(\mathbf{70}_{cs}, 1_c, 2; 8_f), \\
&(\overline{20}_{cs}, 1_c, 4; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle,
\end{aligned} \tag{39}$$

$$\begin{aligned}
|3, 8_f \otimes \overline{10}_f\rangle_2 &= 0.156|896_{cs}, 1_c, 3; 8_f \otimes \overline{10}_f\rangle - 0.364|280_{cs}, 1_c, 3; 8_f \otimes \overline{10}_f\rangle \\
&+ 0.918|35_{cs}, 1_c, 3; 8_f \otimes \overline{10}_f\rangle = -0.422|(\mathbf{70}_{cs}, 8_c, 4; 8_f), (\overline{20}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle \\
&+ 0.629|(\mathbf{70}_{cs}, 8_c, 2; 8_f), (\overline{20}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle + 0.653|(\mathbf{70}_{cs}, 1_c, 2; 8_f), \\
&(\overline{20}_{cs}, 1_c, 4; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle,
\end{aligned} \tag{40}$$

$$\begin{aligned}
|3, 8_f \otimes \overline{10}_f\rangle_3 = & -0.518|896_{cs}, 1_c, 3; 8_f \otimes \overline{10}_f\rangle + 0.762|280_{cs}, 1_c, 3; 8_f \otimes \overline{10}_f\rangle \\
& + 0.389|35_{cs}, 1_c, 3; 8_f \otimes \overline{10}_f\rangle = -0.595|(\mathbf{70}_{cs}, 8_c, 4; 8_f), (\overline{20}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle \\
& + 0.351|(\mathbf{70}_{cs}, 8_c, 2; 8_f), (\overline{20}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle - 0.723|(\mathbf{70}_{cs}, 1_c, 2; 8_f), \\
& (\overline{20}_{cs}, 1_c, 4; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle
\end{aligned} \tag{41}$$

The colormagnetic energy respectively are 24.634C, -12.007C, 7.373C.

C), For the two spin-2 states  $|189_{cs}, 1_c, 5; 8_f \otimes \overline{10}_f\rangle$  and  $|896_{cs}, 1_c, 5; 8_f \otimes \overline{10}_f\rangle$  ( or  $|(\mathbf{70}_{cs}, 8_c, 4; 8_f), (\overline{\mathbf{20}}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 5; 8_f \otimes \overline{10}_f\rangle$ ,  $|(\mathbf{70}_{cs}, 1_c, 2; 8_f), (\overline{\mathbf{20}}_{cs}, 1_c, 4; \overline{10}_f), 1_c, 5; 8_f \otimes \overline{10}_f\rangle$ ), similar to calculating the mixing of the 3 spin-1 states in the above, we can obtain the following eigenstates,

$$\begin{aligned} |5; 8_f \otimes \overline{10}_f\rangle_1 &= \frac{3}{5}|896_{cs}, 1_c, 5; 8_f \otimes \overline{10}_f\rangle + \frac{4}{5}|189_{cs}, 1_c, 5; 8_f \otimes \overline{10}_f\rangle \\ &= -0.447|(\mathbf{70}_{cs}, 8_c, 4; 8_f), (\overline{\mathbf{20}}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 5; 8_f \otimes \overline{10}_f\rangle + 0.894|(\mathbf{70}_{cs}, 1_c, 2; 8_f), \\ &(\overline{\mathbf{20}}_{cs}, 1_c, 4; \overline{10}_f), 1_c, 5; 8_f \otimes \overline{10}_f\rangle, \end{aligned} \quad (42)$$

$$\begin{aligned} |5; 8_f \otimes \overline{10}_f\rangle_2 &= -\frac{4}{5}|896_{cs}, 1_c, 5; 8_f \otimes \overline{10}_f\rangle + \frac{3}{5}|189_{cs}, 1_c, 5; 8_f \otimes \overline{10}_f\rangle \\ &= -0.894|(\mathbf{70}_{cs}, 8_c, 4; 8_f), (\overline{\mathbf{20}}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 5; 8_f \otimes \overline{10}_f\rangle - 0.447|(\mathbf{70}_{cs}, 1_c, 2; 8_f), \\ &(\overline{\mathbf{20}}_{cs}, 1_c, 4; \overline{10}_f), 1_c, 5; 8_f \otimes \overline{10}_f\rangle \end{aligned} \quad (43)$$

The colormagnetic energy are respectively -4C, 16C. The wavefunction for  $\mathbf{20}_{cs} \otimes \overline{\mathbf{70}}_{cs}$  can be obtained by change quarks and antiquarks in the wavefunction Eq.(37), Eq.(39-43), and the corresponding colormagnetic energy is the same.

#### 5. $\mathbf{70}_{cs} \otimes \overline{\mathbf{56}}_{cs}$ and $\mathbf{56}_{cs} \otimes \overline{\mathbf{70}}_{cs}$ .

The multiplets are related by charge conjugation, so we only discuss  $\mathbf{70}_{cs} \otimes \overline{\mathbf{56}}_{cs}$  and the results for  $\mathbf{56}_{cs} \otimes \overline{\mathbf{70}}_{cs}$  can be obtained by exchange quarks and antiquarks. Its flavor representation is  $\mathbf{8}_f \otimes \mathbf{1}_f = \mathbf{8}_f$ , The calculation is analogous to that of  $\mathbf{20}_{cs} \otimes \overline{\mathbf{20}}_{cs}$ , and

$$\mathbf{70}_{cs} \otimes \overline{\mathbf{56}}_{cs} = \mathbf{3200}_{cs} \oplus \mathbf{405}_{cs} \oplus \overline{\mathbf{280}}_{cs} \oplus \mathbf{35}_{cs} \quad (44)$$

with color singlets as followings from Table VII and Table VIII,

$$\begin{aligned} (\mathbf{1}_c, \mathbf{3}) &\subset \mathbf{35}_{cs} \\ (\mathbf{1}_c, \mathbf{3}) &\subset \overline{\mathbf{280}}_{cs} \\ (\mathbf{1}_c, \mathbf{1}) \text{ and } (\mathbf{1}_c, \mathbf{5}) &\subset \mathbf{405}_{cs} \\ (\mathbf{1}_c, \mathbf{3}) \text{ and } (\mathbf{1}_c, \mathbf{5}) &\subset \mathbf{3200}_{cs} \end{aligned} \quad (45)$$

We study each spin states respectively:

A), The spin-0 state is simple, Its wave function is

$$|\mathbf{405}_{cs}, 1_c, 1; 8_f \otimes \mathbf{1}_f\rangle = |(\mathbf{70}_{cs}, 8_c, 2; 8_f), (\overline{\mathbf{56}}_{cs}, 8_c, 2; \mathbf{1}_f), 1_c, 1; 8_f \otimes \mathbf{1}_f\rangle \quad (46)$$

The colormagnetic energy is  $-24C$ .

B), The three spin-1 states  $|35_{cs}, 1_c, 3; 8_f \otimes 1_f\rangle$ ,  $|\overline{280}_{cs}, 1_c, 3; 8_f \otimes 1_f\rangle$  and  $|3200_{cs}, 1_c, 3; 8_f \otimes 1_f\rangle$  ( or  $|(\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; 8_f), (\overline{\mathbf{56}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{4}; 1_f), 1_c, 3; 8_f \otimes 1_f\rangle$ ,  $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; 8_f), (\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; 1_f), 1_c, 3; 8_f \otimes 1_f\rangle$ ,  $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; 8_f), (\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; 1_f), 1_c, 3; 8_f \otimes 1_f\rangle$ ) are mixed by the interaction  $H'$ , and the matrix form of  $H'$  is Eq.(B3). The eigenstates are

$$\begin{aligned} |3, 8_f \otimes 1_f\rangle_1 &= 0.416|(\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; 8_f), (\overline{\mathbf{56}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{4}; 1_f), 1_c, 3; 8_f \otimes 1_f\rangle \\ &- 0.670|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; 8_f), (\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; 1_f), 1_c, 3; 8_f \otimes 1_f\rangle \\ &- 0.615|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; 8_f), (\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; 1_f), 1_c, 3; 8_f \otimes 1_f\rangle, \end{aligned} \quad (47)$$

$$\begin{aligned} |3, 8_f \otimes 1_f\rangle_2 &= 0.732|(\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; 8_f), (\overline{\mathbf{56}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{4}; 1_f), 1_c, 3; 8_f \otimes 1_f\rangle \\ &+ 0.648|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; 8_f), (\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; 1_f), 1_c, 3; 8_f \otimes 1_f\rangle \\ &- 0.212|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; 8_f), (\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; 1_f), 1_c, 3; 8_f \otimes 1_f\rangle, \end{aligned} \quad (48)$$

$$\begin{aligned} |3, 8_f \otimes 1_f\rangle_3 &= 0.540|(\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; 8_f), (\overline{\mathbf{56}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{4}; 1_f), 1_c, 3; 8_f \otimes 1_f\rangle \\ &- 0.362|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; 8_f), (\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; 1_f), 1_c, 3; 8_f \otimes 1_f\rangle \\ &+ 0.760|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; 8_f), (\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; 1_f), 1_c, 3; 8_f \otimes 1_f\rangle \end{aligned} \quad (49)$$

The corresponding eigenvalues are  $-45.078C$ ,  $-14.477C$ ,  $7.555C$ .

C), The two spin-2 states  $|405_{cs}, 1_c, 5; 8_f \otimes 1_f\rangle$  and  $|3200_{cs}, 1_c, 5; 8_f \otimes 1_f\rangle$  are linear combinations of the states  $|(\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; 8_f), (\overline{\mathbf{56}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{4}; 1_f), 1_c, 5; 8_f \otimes 1_f\rangle$  and  $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; 8_f), (\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; 1_f), 1_c, 5; 8_f \otimes 1_f\rangle$ .  $H'$ 's matrix form is Eq.(B4), and the eigenstates are

$$\begin{aligned} |5, 8_f \otimes 1_f\rangle_1 &= 0.845|(\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; 8_f), (\overline{\mathbf{56}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{4}; 1_f), 1_c, 5; 8_f \otimes 1_f\rangle \\ &+ 0.535|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; 8_f), (\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; 1_f), 1_c, 5; 8_f \otimes 1_f\rangle, \end{aligned} \quad (50)$$

$$\begin{aligned} |5, 8_f \otimes 1_f\rangle_2 &= -0.535|(\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; 8_f), (\overline{\mathbf{56}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{4}; 1_f), 1_c, 5; 8_f \otimes 1_f\rangle \\ &+ 0.845|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; 8_f), (\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; 1_f), 1_c, 5; 8_f \otimes 1_f\rangle \end{aligned} \quad (51)$$

The corresponding eigenvalues are  $16C$  and  $-12C$ . The eigenstates for  $\mathbf{56}_{cs} \otimes \overline{\mathbf{70}}_{cs}$  can



be obtained by interchanging the quarks and the antiquarks, and the corresponding eigenvalues are the same as those of  $\mathbf{70}_{cs} \otimes \overline{\mathbf{56}}_{cs}$ .

6.  $\mathbf{70}_{cs} \otimes \overline{\mathbf{70}}_{cs}$ .

Since  $\mathbf{8}_f \otimes \mathbf{8}_f = \mathbf{1}_f \oplus \mathbf{8}_f \oplus \mathbf{8}_f \oplus \mathbf{10}_f \oplus \overline{\mathbf{10}}_f \oplus \mathbf{27}_f$ , the multiplets can be in flavor  $\mathbf{1}_f$ -plet,  $\mathbf{8}_f$ -plet,  $\mathbf{10}_f$ -plet,  $\overline{\mathbf{10}}_f$ -plet or  $\mathbf{27}_f$ -plet. From Fig.1(I) we can see

$$\mathbf{70}_{cs} \otimes \overline{\mathbf{70}}_{cs} = \mathbf{3675}_{cs} \oplus \mathbf{405}_{cs} \oplus \mathbf{280}_{cs} \oplus \overline{\mathbf{280}}_{cs} \oplus \mathbf{35}_{cs} \oplus \mathbf{189}_{cs} \oplus \mathbf{35}_{cs} \oplus \mathbf{1}_{cs} \quad (52)$$

The  $\text{SU}(6)_{cs}$  decomposition with respect to  $\text{SU}(3)_c \otimes \text{SU}(2)_s$  which are listed in Table VII and Table VIII shows the following color singlet,

$$\begin{aligned} (\mathbf{1}_c, \mathbf{1}) &\subset \mathbf{1}_{cs} \\ (\mathbf{1}_c, \mathbf{3}) &\subset \mathbf{35}_{cs} \quad (\text{there exist two } \mathbf{35}_{cs} - \text{plets}) \\ (\mathbf{1}_c, \mathbf{1}) \text{ and } (\mathbf{1}_c, \mathbf{5}) &\subset \mathbf{189}_{cs} \\ (\mathbf{1}_c, \mathbf{3}) &\subset \mathbf{280}_{cs} \\ (\mathbf{1}_c, \mathbf{3}) &\subset \overline{\mathbf{280}}_{cs} \\ (\mathbf{1}_c, \mathbf{1}) \text{ and } (\mathbf{1}_c, \mathbf{5}) &\subset \mathbf{405}_{cs} \\ (\mathbf{1}_c, \mathbf{1}), 2(\mathbf{1}_c, \mathbf{3}), (\mathbf{1}_c, \mathbf{5}) \text{ and } (\mathbf{1}_c, \mathbf{7}) &\subset \mathbf{3675}_{cs} \end{aligned} \quad (53)$$

A), the spin-3 eigenstates, since the  $\text{SU}_c(3) \otimes \text{SU}_s(2)$  decomposition of  $\mathbf{70}_{cs}$  is  $\mathbf{70}_{cs} = (\mathbf{10}_c, \mathbf{2}) + (\mathbf{8}_c, \mathbf{4}) + (\mathbf{8}_c, \mathbf{2}) + (\mathbf{1}_c, \mathbf{2})$ , it is straightforward to get the wave function of the system with spin-3 :

$$|\mathbf{3675}_{cs}, \mathbf{1}_c, \mathbf{7}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle = |(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{7}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle. \quad (54)$$

By the Eq.(2), the corresponding colormagnetic energy is  $16C$ .

B), the spin-2 eigenstates: The three spin-2 states  $|\mathbf{3675}_{cs}, \mathbf{1}_c, \mathbf{5}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ ,  $|\mathbf{405}_{cs}, \mathbf{1}_c, \mathbf{5}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ , and  $|\mathbf{189}_{cs}, \mathbf{1}_c, \mathbf{5}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$  are linear combinations of  $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{5}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ ,  $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{5}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle$  and  $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{5}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ , but they are mixed by the colormagnetic interaction hamiltonian  $H'$ , and  $H'$  can be expressed as the matrix Eq.(B5) in the subspace expanded by the latter three states,

diagonalizing this matrix, we can obtain the eigenstates and eigenvalues. The eigenstates are

$$\begin{aligned} |5, 8_f \otimes 8_f\rangle_1 = & -0.707|(\mathbf{70}_{cs}, 8_c, 4; 8_f), (\overline{\mathbf{70}}_{cs}, 8_c, 2; 8_f), 1_c, 5, 8_f \otimes 8_f\rangle \\ & +0.707|(\mathbf{70}_{cs}, 8_c, 2; 8_f), (\overline{\mathbf{70}}_{cs}, 8_c, 4; 8_f), 1_c, 5, 8_f \otimes 8_f\rangle, \end{aligned} \quad (55)$$

$$\begin{aligned} |5, 8_f \otimes 8_f\rangle_2 = & -0.526|(\mathbf{70}_{cs}, 8_c, 4; 8_f), (\overline{\mathbf{70}}_{cs}, 8_c, 4; 8_f), 1_c, 5, 8_f \otimes 8_f\rangle \\ & -0.602|(\mathbf{70}_{cs}, 8_c, 4; 8_f), (\overline{\mathbf{70}}_{cs}, 8_c, 2; 8_f), 1_c, 5, 8_f \otimes 8_f\rangle \\ & -0.602|(\mathbf{70}_{cs}, 8_c, 2; 8_f), (\overline{\mathbf{70}}_{cs}, 8_c, 4; 8_f), 1_c, 5, 8_f \otimes 8_f\rangle, \end{aligned} \quad (56)$$

$$\begin{aligned} |5, 8_f \otimes 8_f\rangle_3 = & 0.851|(\mathbf{70}_{cs}, 8_c, 4; 8_f), (\overline{\mathbf{70}}_{cs}, 8_c, 4; 8_f), 1_c, 5, 8_f \otimes 8_f\rangle \\ & -0.372|(\mathbf{70}_{cs}, 8_c, 4; 8_f), (\overline{\mathbf{70}}_{cs}, 8_c, 2; 8_f), 1_c, 5, 8_f \otimes 8_f\rangle \\ & -0.372|(\mathbf{70}_{cs}, 8_c, 2; 8_f), (\overline{\mathbf{70}}_{cs}, 8_c, 4; 8_f), 1_c, 5, 8_f \otimes 8_f\rangle \end{aligned} \quad (57)$$

And the corresponding eigenvalues are  $16C$ ,  $-12.944C$ ,  $4.944C$ .

C), the spin-1 eigenstates: The six spin-1 states  $|\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; 8_f \otimes 8_f\rangle_1$ ,  $|\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; 8_f \otimes 8_f\rangle_2$ ,  $|\mathbf{280}_{cs}, \mathbf{1}_c, \mathbf{3}; 8_f \otimes 8_f\rangle$ ,  $|\mathbf{280}_{cs}, \mathbf{1}_c, \mathbf{3}; 8_f \otimes 8_f\rangle$ ,  $|\mathbf{3675}_{cs}, \mathbf{1}_c, \mathbf{3}; 8_f \otimes 8_f\rangle_1$  and  $|\mathbf{3675}_{cs}, \mathbf{1}_c, \mathbf{3}; 8_f \otimes 8_f\rangle_2$  are linear combination of the six direct product states  $|(\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; 8_f), (\overline{\mathbf{70}}_{cs}, \mathbf{10}_c, \mathbf{2}; 8_f), 1_c, \mathbf{3}; 8_f \otimes 8_f\rangle$ ,  $|(\mathbf{70}_{cs}, 8_c, 4; 8_f), (\overline{\mathbf{70}}_{cs}, 8_c, 4; 8_f), 1_c, \mathbf{3}; 8_f \otimes 8_f\rangle$ ,  $|(\mathbf{70}_{cs}, 8_c, 4; 8_f), (\overline{\mathbf{70}}_{cs}, 8_c, 2; 8_f), 1_c, \mathbf{3}; 8_f \otimes 8_f\rangle$ ,  $|(\mathbf{70}_{cs}, 8_c, 2; 8_f), (\overline{\mathbf{70}}_{cs}, 8_c, 4; 8_f), 1_c, \mathbf{3}; 8_f \otimes 8_f\rangle$ ,  $|(\mathbf{70}_{cs}, 8_c, 2; 8_f), (\overline{\mathbf{70}}_{cs}, 8_c, 2; 8_f), 1_c, \mathbf{3}; 8_f \otimes 8_f\rangle$  and  $|(\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; 8_f), (\overline{\mathbf{70}}_{cs}, \mathbf{1}_c, \mathbf{2}; 8_f), 1_c, \mathbf{3}; 8_f \otimes 8_f\rangle$ . Noting that there are two  $\mathbf{35}_{cs}$  in the outer product of  $\mathbf{70}_{cs} \otimes \overline{\mathbf{70}}_{cs}$ , the states  $|\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; 8_f \otimes 8_f\rangle_1$  and  $|\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; 8_f \otimes 8_f\rangle_2$  represent the two states respectively belonging to the two  $\mathbf{35}_{cs}$ .  $(\mathbf{1}_c, \mathbf{3})$  appears twice in the decomposition of  $\mathbf{3675}_{cs}$ , then the states  $|\mathbf{3675}_{cs}, \mathbf{1}_c, \mathbf{3}; 8_f \otimes 8_f\rangle_1$  and  $|\mathbf{3675}_{cs}, \mathbf{1}_c, \mathbf{3}; 8_f \otimes 8_f\rangle_2$  denote the two states corresponding to the two  $(\mathbf{1}_c, \mathbf{3})$  respectively. Once again the states of definite spin and  $\text{SU}(6)_{cs}$  representation are mixed by  $H'$ . By using the matrix representation of  $H'$ , i.e., Eq.(B6), the eigenstates are

$$|\mathbf{3}, 8_f \otimes 8_f\rangle_1 = 0.2|(\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; 8_f), (\overline{\mathbf{70}}_{cs}, \mathbf{10}_c, \mathbf{2}; 8_f), 1_c, \mathbf{3}; 8_f \otimes 8_f\rangle$$



$$\begin{aligned}
& -0.239|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle \\
& -0.239|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle \\
& -0.649|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle \\
& -0.441|(\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle,
\end{aligned} \tag{62}$$

$$\begin{aligned}
& |\mathbf{3}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle_6 = -0.707|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle \\
& +0.707|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle
\end{aligned} \tag{63}$$

Their corresponding eigenvalues are  $-29.461C$ ,  $27.631C$ ,  $-20.293C$ ,  $-10.667C$ ,  $3.456C$ ,  $-2.667C$  respectively.

D), the spin-0 eigenstates: similarly the four spin-0 states  $|\mathbf{1}_{cs}, \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ ,  $|\mathbf{189}_{cs}, \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ ,  $|\mathbf{405}_{cs}, \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ ,  $|\mathbf{3675}_{cs}, \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ , are linear combination of the four states  $|(\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ ,  $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ ,  $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$  and  $|(\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ .  $H'$ 's matrix form is Eq.(B7), thus the eigenstates are

$$\begin{aligned}
& |\mathbf{1}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle_1 = 0.394|(\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle \\
& +0.604|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle \\
& -0.576|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle \\
& -0.384|(\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle,
\end{aligned} \tag{64}$$

$$\begin{aligned}
& |\mathbf{1}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle_2 = -0.725|(\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle \\
& -0.649|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle \\
& +0.229|(\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle,
\end{aligned} \tag{65}$$

$$\begin{aligned}
& |\mathbf{1}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle_3 = 0.371|(\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle \\
& +0.232|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle \\
& -0.099|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle \\
& +0.894|(\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle,
\end{aligned} \tag{66}$$

$$\begin{aligned}
|1, 8_f \otimes 8_f\rangle_4 &= 0.425|(\mathbf{70}_{cs}, \mathbf{10}_c, 2; 8_f), (\overline{\mathbf{70}}_{cs}, \overline{\mathbf{10}}_c, 2; 8_f), 1_c, 1; 8_f \otimes 8_f\rangle \\
&- 0.763|(\mathbf{70}_{cs}, 8_c, 4; 8_f), (\overline{\mathbf{70}}_{cs}, 8_c, 4; 8_f), 1_c, 1; 8_f \otimes 8_f\rangle \\
&- 0.487|(\mathbf{70}_{cs}, 8_c, 2; 8_f), (\overline{\mathbf{70}}_{cs}, 8_c, 2; 8_f), 1_c, 1; 8_f \otimes 8_f\rangle \\
&- 0.033|(\mathbf{70}_{cs}, 1_c, 2; 8_f), (\overline{\mathbf{70}}_{cs}, 1_c, 2; 8_f), 1_c, 1; 8_f \otimes 8_f\rangle
\end{aligned} \tag{67}$$

The eigenvalues are  $-45.547C$ ,  $16C$ ,  $-12.671C$ ,  $2.218C$  respectively.

Finally, as a matter of convenience, we list all the QCD-allowed states (i.e., color singlets) and their colormagnetic energy in the Table I. From this Table, we learned at first that the configurations corresponding to the  $\Lambda\overline{\Lambda}$ ,  $p\overline{\Lambda}$ ,  $p\overline{p}$ ,  $\dots$  enhancements can have rather large negative colormagnetic energy. This indicates that these states should be visible in experiments, and therefore these results are consistent with the experiments reported in Ref.[2-8]. Secondly, we can see also that the binding color-magnetic energy of spin-0 and spin-1 states are larger than ones of the states with spin-2 and spin-3.

Specifically, and more concretely,  $|1, 1_f \otimes 1_f\rangle_1$  (the wave function is Eq.(31)) has the largest colormagnetic energy  $-82.533C$ , and it has the same quantum numbers with the  $\Lambda\overline{\Lambda}$  enhancement [8], so this state possibly is just the candidate  $\Lambda\overline{\Lambda}$  enhancement. And the state  $|3, 1_f \otimes 1_f\rangle_1$  with the wave function Eq.(33) has colormagnetic energy  $-48.331C$ , then this state is also rather stable. This spin-1  $\Lambda\overline{\Lambda}$  enhancement as a prediction is expected to be observed by BES, Bell or CLEO in future. By similar analysis, the colormagnetic interaction energy of the state  $|1, 8_f \otimes 8_f\rangle_1$  (the wave function is Eq.(64)) is  $-45.547C$ , and the  $p\overline{p}$  enhancement[2, 4] should belong to these states. Since the colormagnetic energy of the proton or antiproton is  $-8C$  (please see Table II), we can estimate that the mass defect of the  $p\overline{p}$  enhancement approximately is  $29.547C$  which is compatible with the experimental observation, that the  $p\overline{p}$  enhancement is bellow the threshold. Also we see that the states  $|3, 8_f \otimes 8_f\rangle_1$  whose wave function is Eq.(58) has colormagnetic energy  $-29.461C$ , so these states could also be seen experimentally. Therefore we conjecture that both the spin-0 and spin-1  $p\overline{p}$  enhancement may have been observed by BES[2, 4]. The states  $|1, 8_f \otimes 1_f\rangle$  (the wave function is given by Eq.(46)) has colormagnetic energy  $-24C$ , and the  $p\overline{\Lambda}$  enhancement[5, 7] can belong to these states. However the states  $|3, 8_f \otimes 1_f\rangle_1$  whose wave function is given

by Eq.(47) has colormagnetic energy -45.078C, which is larger than the colormagnetic energy of the state  $|\mathbf{1}, \mathbf{8}_f \otimes \mathbf{1}_f\rangle$ , so it indicates that the vector  $p\bar{\Lambda}$  enhancement may be favored over the scalar one. Because the  $p\bar{\Lambda}$  enhancement has been seen in the process  $J/\psi \rightarrow p\bar{\Lambda}K^-$  and  $\psi' \rightarrow p\bar{\Lambda}K^-$ , and  $J/\psi$  and  $\psi'$  are vector mesons,  $K^-$  is a pseudoscalar meson, the relative angular momentum between  $K^-$  and the  $p\bar{\Lambda}$  enhancement  $l = 0$  is favored, if the  $p\bar{\Lambda}$  enhancement is a vector state. We also conjecture that both the spin-0 and spin-1  $p\bar{\Lambda}$  enhancement has been observed by BES but the main content is spin-1  $p\bar{\Lambda}$  enhancement.

#### IV. THE INTERESTING CONFIGURATIONS

Quark correlation turns out to play important role in multiquark states [15, 16], so we consider the effect of quark correlation in the  $q^3\bar{q}^3$  system. Generally in the  $q^3\bar{q}^3$  system there are many possible configuration:  $(q) - (q^2\bar{q}^3)$ ,  $(\bar{q}) - (q^3\bar{q}^2)$ ,  $(q^2) - (q\bar{q}^3)$ ,  $(q\bar{q}) - (q^2\bar{q}^2)$ ,  $(\bar{q}^2) - (q^3\bar{q})$ ,  $(q^3) - (\bar{q}^3)$ ,  $(q^2\bar{q}) - (q\bar{q}^2)$  and so on. In this paper, we only consider the symmetric configurations:  $(q^3) - (\bar{q}^3)$ ,  $(q^2\bar{q}) - (q\bar{q}^2)$ , where under the changes  $q \leftrightarrow \bar{q}$ , the 1<sup>st</sup>-cluster and 2<sup>nd</sup>-cluster interchange each other. The reasons why we choose these configuration are both because triquark correlation has been proved to be important in pentaquark[15], and because the configurations are more symmetric than others. The cluster structure is postulated instead of being established by a dynamical calculation. But with the development of lattice QCD, we could study which configurations are more stable than others through lattice numerical calculations[22]. It is assumed the two clusters are separated by a distance larger than the range of the colormagnetic force which is of short range and are kept together by the colorelectric force. Consequently, the colormagnetic hyperfine interaction operates only within each cluster, but is not felt between them.

Each cluster can be in a variety of  $SU(3)_c$  representations, and eventually two clusters together form the color singlet physical baryonium state. However, if two clusters are color singlets respectively, they will interact only via the van der Waal's force of QCD and will dissociate immediately into a baryon and a antibaryon, or be bound very weakly to form a very unstable particle which can not be regarded as a narrow resonance of baronium that responds to the enhancement.

If the two clusters are connected via the color triplet bonds, the creation of a  $q\bar{q}$  pair

TABLE I: Colormagnetic interaction energy of  $q^3\bar{q}^3$  eigenstates in common quark model, and the conjugates states are not included.

| State                                                             | Wave Function | Colormagnetic<br>Interaction Energy | State                                                             | Wave Function | Colormagnetic<br>Interaction Energy |
|-------------------------------------------------------------------|---------------|-------------------------------------|-------------------------------------------------------------------|---------------|-------------------------------------|
| $ \mathbf{1}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle_1$         | Eq.(64)       | -45.547C                            | $ \mathbf{1}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle_2$         | Eq.(65)       | 16C                                 |
| $ \mathbf{1}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle_3$         | Eq.(66)       | -12.671C                            | $ \mathbf{1}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle_4$         | Eq.(67)       | 2.218C                              |
| $ \mathbf{1}, \mathbf{8}_f \otimes \mathbf{1}_f\rangle$           | Eq.(46)       | -24C                                | $ \mathbf{1}, \mathbf{8}_f \otimes \bar{\mathbf{10}}_f\rangle$    | Eq.(37)       | 8C                                  |
| $ \mathbf{1}, \mathbf{1}_f \otimes \bar{\mathbf{10}}_f\rangle$    | Eq.(25)       | 16C                                 | $ \mathbf{1}, \mathbf{1}_f \otimes \mathbf{1}_f\rangle_1$         | Eq.(31)       | -82.533C                            |
| $ \mathbf{1}, \mathbf{1}_f \otimes \mathbf{1}_f\rangle_2$         | Eq.(32)       | -29.467C                            | $ \mathbf{1}, \mathbf{10}_f \otimes \bar{\mathbf{10}}_f\rangle_1$ | Eq.(21)       | 25.889C                             |
| $ \mathbf{1}, \mathbf{10}_f \otimes \bar{\mathbf{10}}_f\rangle_2$ | Eq.(22)       | -9.889C                             | $ \mathbf{3}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle_1$         | Eq.(58)       | -29.461C                            |
| $ \mathbf{3}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle_2$         | Eq.(59)       | 27.631C                             | $ \mathbf{3}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle_3$         | Eq.(60)       | -20.293C                            |
| $ \mathbf{3}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle_4$         | Eq.(61)       | -10.667C                            | $ \mathbf{3}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle_5$         | Eq.(62)       | 3.456C                              |
| $ \mathbf{3}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle_6$         | Eq.(63)       | -2.667C                             | $ \mathbf{3}, \mathbf{8}_f \otimes \mathbf{1}_f\rangle_1$         | Eq.(47)       | -45.078C                            |
| $ \mathbf{3}, \mathbf{8}_f \otimes \mathbf{1}_f\rangle_2$         | Eq.(48)       | -14.477C                            | $ \mathbf{3}, \mathbf{8}_f \otimes \mathbf{1}_f\rangle_3$         | Eq.(49)       | 7.555C                              |
| $ \mathbf{3}, \mathbf{8}_f \otimes \bar{\mathbf{10}}_f\rangle_1$  | Eq.(39)       | 24.634C                             | $ \mathbf{3}, \mathbf{8}_f \otimes \bar{\mathbf{10}}_f\rangle_2$  | Eq.(40)       | -12.007C                            |
| $ \mathbf{3}, \mathbf{8}_f \otimes \bar{\mathbf{10}}_f\rangle_3$  | Eq.(41)       | 7.373C                              | $ \mathbf{3}, \mathbf{1}_f \otimes \bar{\mathbf{10}}_f\rangle$    | Eq.(26)       | $-\frac{32}{3}C$                    |
| $ \mathbf{3}, \mathbf{1}_f \otimes \mathbf{1}_f\rangle_1$         | Eq.(33)       | -48.331C                            | $ \mathbf{3}, \mathbf{1}_f \otimes \mathbf{1}_f\rangle_2$         | Eq.(34)       | -5.003C                             |
| $ \mathbf{3}, \mathbf{10}_f \otimes \bar{\mathbf{10}}_f\rangle_1$ | Eq.(19)       | 34.397C                             | $ \mathbf{3}, \mathbf{10}_f \otimes \bar{\mathbf{10}}_f\rangle_2$ | Eq.(20)       | 8.269C                              |
| $ \mathbf{5}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle_1$         | Eq.(55)       | 16C                                 | $ \mathbf{5}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle_2$         | Eq.(56)       | -12.944C                            |
| $ \mathbf{5}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle_3$         | Eq.(57)       | 4.944C                              | $ \mathbf{5}, \mathbf{8}_f \otimes \mathbf{1}_f\rangle_1$         | Eq.(50)       | 16C                                 |
| $ \mathbf{5}, \mathbf{8}_f \otimes \mathbf{1}_f\rangle_2$         | Eq.(51)       | -12C                                | $ \mathbf{5}, \mathbf{8}_f \otimes \bar{\mathbf{10}}_f\rangle_1$  | Eq.(42)       | -4C                                 |
| $ \mathbf{5}, \mathbf{8}_f \otimes \bar{\mathbf{10}}_f\rangle_2$  | Eq.(43)       | 16C                                 | $ \mathbf{5}, \mathbf{1}_f \otimes \mathbf{1}_f\rangle$           | Eq.(30)       | -16C                                |
| $ \mathbf{5}, \mathbf{10}_f \otimes \bar{\mathbf{10}}_f\rangle$   | Eq.(16)       | 16C                                 | $ \mathbf{7}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle$           | Eq.(54)       | 16C                                 |
| $ \mathbf{7}, \mathbf{1}_f \otimes \mathbf{1}_f\rangle$           | Eq.(29)       | 16C                                 | $ \mathbf{7}, \mathbf{10}_f \otimes \bar{\mathbf{10}}_f\rangle$   | Eq.(15)       | 16C                                 |

permits the system to break up into two separate states which are color singlet. So the lifetime of the state is comparable to that of the excited  $q\bar{q}$  meson states or excited  $qqq$  baryon states, and the decay width is larger than 100 MeV, since the ordinary meson or baryon states are also bound together by color triplet bond[17, 19]. In Jaffe's language[19]

the process is superallowed if it is heavy enough. Therefore, for our purpose, we only consider the configurations that the two clusters are separately in higher dimensional  $SU(3)_c$  representation, such as the  $\mathbf{15}_c - \overline{\mathbf{15}}_c, \mathbf{10}_c - \overline{\mathbf{10}}_c, \mathbf{8}_c - \mathbf{8}_c, \mathbf{6}_c - \overline{\mathbf{6}}_c$  configurations.

### A. The $(q^3) - (\bar{q}^3)$ Configuration

The configuration is schematically shown in Fig.4, from Eq.(11) we see that the allowed states of the  $q^3$  cluster are

$$|\mathbf{70}_{cs}, \mathbf{8}_f\rangle, |\mathbf{50}_{cs}, \mathbf{1}_f\rangle, |\mathbf{20}_{cs}, \mathbf{10}_f\rangle \quad (68)$$

The  $SU(6)_{cs}$  irreducible representation  $\mathbf{70}_{cs}, \mathbf{56}_{cs}, \mathbf{20}_{cs}$  are decomposed as follows under  $SU(3)_c \otimes SU(2)_s$

$$\begin{aligned} \mathbf{70}_{cs} &= (\mathbf{10}_c, \mathbf{2}) + (\mathbf{8}_c, \mathbf{4}) + (\mathbf{8}_c, \mathbf{2}) + (\mathbf{1}_c, \mathbf{2}), \\ \mathbf{56}_{cs} &= (\mathbf{10}_c, \mathbf{4}) + (\mathbf{8}_c, \mathbf{2}), \\ \mathbf{20}_{cs} &= (\mathbf{8}_c, \mathbf{2}) + (\mathbf{1}_c, \mathbf{4}), \end{aligned} \quad (69)$$

where the first and the second numbers in the bracket are respectively the irreducible representations of  $SU(3)_c$  and  $SU(2)_s$ . By using the conventional notation  $|D_{cs}, D_c, 2S+1; D_f\rangle$  to express the states, the possible states of the  $q^3$  cluster are

$$\begin{aligned} &|\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; \mathbf{8}_f\rangle, |\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f\rangle, |\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle, |\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f\rangle, |\mathbf{56}_{cs}, \mathbf{10}_c, \mathbf{4}; \mathbf{1}_f\rangle, \\ &|\mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f\rangle, |\mathbf{20}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{10}_f\rangle, |\mathbf{20}_{cs}, \mathbf{1}_c, \mathbf{4}; \mathbf{10}_f\rangle. \end{aligned} \quad (70)$$

Noting the symmetry group of both color and flavor are  $SU(3)$ , we use a subscript c and f to distinguish them. Similarly the possible states of  $\bar{q}^3$  cluster are

$$\begin{aligned} &|\overline{\mathbf{70}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{2}; \mathbf{8}_f\rangle, |\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f\rangle, |\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle, |\overline{\mathbf{70}}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f\rangle, |\overline{\mathbf{56}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{4}; \mathbf{1}_f\rangle, \\ &|\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f\rangle, |\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f\rangle, |\overline{\mathbf{20}}_{cs}, \mathbf{1}_c, \mathbf{4}; \overline{\mathbf{10}}_f\rangle \end{aligned} \quad (71)$$

where  $\mathbf{20}_{cs} = \overline{\mathbf{20}}_{cs}$ .

The above states are eigenstates of the colormagnetic interaction hamiltonian  $H'$ , and we can calculate the corresponding eigenvalues, the results are listed in Table II.

From Eq.(70) we can learn that the color singlet states  $|\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f\rangle$  and  $|\mathbf{20}_{cs}, \mathbf{1}_c, \mathbf{4}; \mathbf{10}_f\rangle$  are the usual baryon octet and baryon decuplet respectively, and from



| States                                                                                                                                                               | Eigenvalue of the Hamiltonian $H'$ |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------|
| $ \mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; \mathbf{8}_f\rangle$ ( or $ \overline{\mathbf{70}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{2}; \mathbf{8}_f\rangle$ ) | $4C$                               |
| $ \mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f\rangle$ ( or $ \overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f\rangle$ )              | $2C$                               |
| $ \mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle$ ( or $ \overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle$ )              | $-2C$                              |
| $ \mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f\rangle$ ( or $ \overline{\mathbf{70}}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f\rangle$ )              | $-8C$                              |
| $ \mathbf{56}_{cs}, \mathbf{10}_c, \mathbf{4}; \mathbf{1}_f\rangle$ ( or $ \overline{\mathbf{56}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{4}; \mathbf{1}_f\rangle$ ) | $-4C$                              |
| $ \mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f\rangle$ ( or $ \overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f\rangle$ )              | $-14C$                             |
| $ \mathbf{20}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{10}_f\rangle$ ( or $ \overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f\rangle$ ) | $10C$                              |
| $ \mathbf{20}_{cs}, \mathbf{1}_c, \mathbf{4}; \mathbf{10}_f\rangle$ ( or $ \overline{\mathbf{20}}_{cs}, \mathbf{1}_c, \mathbf{4}; \overline{\mathbf{10}}_f\rangle$ ) | $8C$                               |

TABLE II: The possible states of the cluster  $q^3$  and the corresponding eigenvalues of the color-magnetic interaction hamilton  $H'$

Table II, their mass difference is  $16C$ , The states of the total system are the product of the subsystem  $q^3$ 's states Eq.(70) and the subsystem  $\bar{q}^3$ 's states Eq.(71). It is obvious that the system can be in the states

$$\begin{aligned}
& |\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f\rangle |\overline{\mathbf{70}}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f\rangle, \quad |\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f\rangle |\overline{\mathbf{20}}_{cs}, \mathbf{1}_c, \mathbf{4}; \overline{\mathbf{10}}_f\rangle \\
& |\mathbf{20}_{cs}, \mathbf{1}_c, \mathbf{4}; \mathbf{10}_f\rangle |\overline{\mathbf{70}}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f\rangle, \quad |\mathbf{20}_{cs}, \mathbf{1}_c, \mathbf{4}; \mathbf{10}_f\rangle |\overline{\mathbf{20}}_{cs}, \mathbf{1}_c, \mathbf{4}; \overline{\mathbf{10}}_f\rangle
\end{aligned} \tag{72}$$

Their colormagnetic energy is  $(-16C, 0, 0, 16C)$ . In the above four states the two clusters are connected by color singlet bond. They easily dissociate into a baryon and an antibaryon just as we mentioned in the above, so we do not discuss these states further. We are interested in the states in which the two clusters are connected by color **10**-plet bond or color **8**-plet bond. It is easy to write out the possible states of this system, to reveal their flavor structure and the spin structure, and to calculate colormagnetic interaction energy. The results are listed in the Table III. We address that the clusters in these states are with color, and can not be observed individually, but the combines of them may be colorless and emerge as baryoniums. Using the results in the Table III, the discussions on enhancement are in order:

1.  $\Lambda_1 \bar{\Lambda}_1$  enhancement: From the Table III, we can see that the state  $|\mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f\rangle |\overline{\mathbf{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f\rangle$  with flavor  $\mathbf{1}_f$  and spin equaling **0** or **1** have the largest colormagnetic energy  $-28C$ , so this quark configuration is rather stable. Since

| States of the Total System                                                                                                                                  | SU <sub>f</sub> (3) Structure                                                                                                    | the Spin Structure                               | the Mass Splitting |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------|--------------------|
| $ \mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; \mathbf{8}_f\rangle  \mathbf{\overline{70}}_{cs}, \mathbf{\overline{10}}_c, \mathbf{2}; \mathbf{8}_f\rangle$ | $\mathbf{1}_f \oplus \mathbf{8}_f \oplus \mathbf{8}_f \oplus \mathbf{10}_f \oplus \mathbf{\overline{10}}_f \oplus \mathbf{27}_f$ | $\mathbf{0}, \mathbf{1}$                         | 8C                 |
| $ \mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; \mathbf{8}_f\rangle  \mathbf{\overline{56}}_{cs}, \mathbf{\overline{10}}_c, \mathbf{4}; \mathbf{1}_f\rangle$ | $\mathbf{8}_f$                                                                                                                   | $\mathbf{1}, \mathbf{2}$                         | 0                  |
| $ \mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f\rangle  \mathbf{\overline{70}}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f\rangle$              | $\mathbf{1}_f \oplus \mathbf{8}_f \oplus \mathbf{8}_f \oplus \mathbf{10}_f \oplus \mathbf{\overline{10}}_f \oplus \mathbf{27}_f$ | $\mathbf{0}, \mathbf{1}, \mathbf{2}, \mathbf{3}$ | 4C                 |
| $ \mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f\rangle  \mathbf{\overline{70}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle$              | $\mathbf{1}_f \oplus \mathbf{8}_f \oplus \mathbf{8}_f \oplus \mathbf{10}_f \oplus \mathbf{\overline{10}}_f \oplus \mathbf{27}_f$ | $\mathbf{1}, \mathbf{2}$                         | 0                  |
| $ \mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f\rangle  \mathbf{\overline{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f\rangle$              | $\mathbf{8}_f$                                                                                                                   | $\mathbf{1}, \mathbf{2}$                         | -12C               |
| $ \mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f\rangle  \mathbf{\overline{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{\overline{10}}_f\rangle$  | $\mathbf{8}_f \oplus \mathbf{\overline{10}}_f \oplus \mathbf{27}_f \oplus \mathbf{\overline{35}}_f$                              | $\mathbf{1}, \mathbf{2}$                         | 12C                |
| $ \mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle  \mathbf{\overline{70}}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f\rangle$              | $\mathbf{1}_f \oplus \mathbf{8}_f \oplus \mathbf{8}_f \oplus \mathbf{10}_f \oplus \mathbf{\overline{10}}_f \oplus \mathbf{27}_f$ | $\mathbf{1}, \mathbf{2}$                         | 0                  |
| $ \mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle  \mathbf{\overline{70}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle$              | $\mathbf{1}_f \oplus \mathbf{8}_f \oplus \mathbf{8}_f \oplus \mathbf{10}_f \oplus \mathbf{\overline{10}}_f \oplus \mathbf{27}_f$ | $\mathbf{0}, \mathbf{1}$                         | -4C                |
| $ \mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle  \mathbf{\overline{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f\rangle$              | $\mathbf{8}_f$                                                                                                                   | $\mathbf{0}, \mathbf{1}$                         | -16C               |
| $ \mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle  \mathbf{\overline{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{\overline{10}}_f\rangle$  | $\mathbf{8}_f \oplus \mathbf{\overline{10}}_f \oplus \mathbf{27}_f \oplus \mathbf{\overline{35}}_f$                              | $\mathbf{0}, \mathbf{1}$                         | 8C                 |
| $ \mathbf{56}_{cs}, \mathbf{10}_c, \mathbf{4}; \mathbf{1}_f\rangle  \mathbf{\overline{70}}_{cs}, \mathbf{\overline{10}}_c, \mathbf{2}; \mathbf{8}_f\rangle$ | $\mathbf{8}_f$                                                                                                                   | $\mathbf{1}, \mathbf{2}$                         | 0                  |
| $ \mathbf{56}_{cs}, \mathbf{10}_c, \mathbf{4}; \mathbf{1}_f\rangle  \mathbf{\overline{56}}_{cs}, \mathbf{\overline{10}}_c, \mathbf{4}; \mathbf{1}_f\rangle$ | $\mathbf{1}_f$                                                                                                                   | $\mathbf{0}, \mathbf{1}, \mathbf{2}, \mathbf{3}$ | -8C                |
| $ \mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f\rangle  \mathbf{\overline{70}}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f\rangle$              | $\mathbf{8}_f$                                                                                                                   | $\mathbf{1}, \mathbf{2}$                         | -12C               |
| $ \mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f\rangle  \mathbf{\overline{70}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle$              | $\mathbf{8}_f$                                                                                                                   | $\mathbf{0}, \mathbf{1}$                         | -16C               |
| $ \mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f\rangle  \mathbf{\overline{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f\rangle$              | $\mathbf{1}_f$                                                                                                                   | $\mathbf{0}, \mathbf{1}$                         | -28C               |
| $ \mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f\rangle  \mathbf{\overline{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{\overline{10}}_f\rangle$  | $\mathbf{\overline{10}}_f$                                                                                                       | $\mathbf{0}, \mathbf{1}$                         | -4C                |
| $ \mathbf{20}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{10}_f\rangle  \mathbf{\overline{70}}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f\rangle$             | $\mathbf{8}_f \oplus \mathbf{10}_f \oplus \mathbf{27}_f \oplus \mathbf{35}_f$                                                    | $\mathbf{1}, \mathbf{2}$                         | 12C                |
| $ \mathbf{20}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{10}_f\rangle  \mathbf{\overline{70}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle$             | $\mathbf{8}_f \oplus \mathbf{10}_f \oplus \mathbf{27}_f \oplus \mathbf{35}_f$                                                    | $\mathbf{0}, \mathbf{1}$                         | 8C                 |
| $ \mathbf{20}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{10}_f\rangle  \mathbf{\overline{56}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f\rangle$             | $\mathbf{10}_f$                                                                                                                  | $\mathbf{0}, \mathbf{1}$                         | -4C                |
| $ \mathbf{20}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{10}_f\rangle  \mathbf{\overline{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{\overline{10}}_f\rangle$ | $\mathbf{1}_f \oplus \mathbf{8}_f \oplus \mathbf{27}_f \oplus \mathbf{64}_f$                                                     | $\mathbf{0}, \mathbf{1}$                         | 20C                |

TABLE III: The allowed states of the total system in the configuration  $(q^3) - (\overline{q}^3)$  and the mass splitting.

the two clusters are all in flavor  $\mathbf{1}_f$ , they are of flavor  $\Lambda_1$ ,  $\overline{\Lambda}_1$  with color octet. It is well known that in the color singlet ground state multiplet, the SU<sub>f</sub>(3) flavor singlet  $\Lambda_1$  is forbidden by Fermi statistics[23]. But the above composite boson state of  $(\Lambda_1 \overline{\Lambda}_1)$  can belong to color singlet, and is not forbidden. And its quantum numbers are same as ones of the state  $(\Lambda \overline{\Lambda})$ . This  $\Lambda_1 \overline{\Lambda}_1$  (or  $\Lambda \overline{\Lambda}$ ) enhancement has been reported in Ref.[8].

2.  $p\bar{\Lambda}_1$  enhancement: Both the states  $|\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle |\mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f\rangle$  and  $|\mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f\rangle |\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle$  have colormagnetic energy  $-16C$ , they are in the flavor  $\mathbf{8}_f$ -plet with spin  $\mathbf{0}$  or  $\mathbf{1}$ . The total system can be in the  $p\bar{\Lambda}$  like state because one of the clusters is in flavor representation  $\mathbf{1}_f$ . One of these states has also been reported by BES[5] and Belle[7], but its spin can not be fixed.
3.  $p\bar{p}$  enhancement: It is obvious that the  $p\bar{p}$  enhancement[2, 4] possibly belongs to the states  $|\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; \mathbf{8}_f\rangle |\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; \mathbf{8}_f\rangle$ ,  $|\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f\rangle |\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f\rangle$ ,  $|\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle |\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle$ ,  $|\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle |\mathbf{20}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{10}_f\rangle$ ,  $|\mathbf{20}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{10}_f\rangle |\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle$ , and the corresponding colormagnetic energy are  $8C$ ,  $4C$ ,  $-4C$ ,  $8C$ ,  $8C$  respectively. This indicates that among them, the quark configuration  $|\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle |\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle$  is more stable. Namely, in the  $(q^3) - (\bar{q}^3)$  configuration, the  $p\bar{p}$  enhancement most possibly belongs to the states  $|\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle |\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f\rangle$ .

Noting that both spin-0, spin-1  $p\bar{p}$  enhancements and spin-0, spin-1  $p\bar{\Lambda}$  enhancements are predicted.

## B. The $(q^2\bar{q}) - (q\bar{q}^2)$ Configuration

The configuration is schematically shown in Fig.5. Due to Fermi statistics, the two quarks in the cluster belong to the  $(\mathbf{6}_{cs} \otimes \mathbf{6}_{cs})_S = \mathbf{21}_{cs}$  dimensional representation of  $SU(6)_{cs}$  and irreducible  $SU(3)_f$  representation  $\mathbf{3}_f$ , or to  $(\mathbf{6}_{cs} \otimes \mathbf{6}_{cs})_A = \mathbf{15}_{cs}$  dimensional representation of  $SU(6)_{cs}$  and  $SU(3)_f$  representation  $\mathbf{6}_f$ . Performing the outer product of  $SU(6)_{cs}$  representations

$$\mathbf{21}_{cs} \otimes \mathbf{6}_{cs} = \mathbf{6}_{cs} \oplus \mathbf{120}_{cs}, \quad \mathbf{15}_{cs} \otimes \mathbf{6}_{cs} = \mathbf{6}_{cs} \oplus \mathbf{84}_{cs}, \quad (73)$$

we find out that the cluster  $q^2\bar{q}$  can be in the  $SU(6)_{cs}$  representation  $\mathbf{6}_{cs}$ ,  $\mathbf{84}_{cs}$  or  $\mathbf{120}_{cs}$ , and the corresponding  $SU(3)_c \otimes SU(2)_s$  decompositions are as followings

$$\begin{aligned} \mathbf{6}_{cs} &= (\mathbf{3}_c, \mathbf{2}), \quad \mathbf{15}_{cs} = (\mathbf{6}_c, \mathbf{1}) + (\mathbf{3}_c, \mathbf{3}), \quad \mathbf{21}_{cs} = (\mathbf{6}_c, \mathbf{3}) + (\mathbf{3}_c, \mathbf{1}) \\ \mathbf{84}_{cs} &= (\mathbf{3}_c, \mathbf{2}) + (\mathbf{3}_c, \mathbf{4}) + (\mathbf{6}_c, \mathbf{2}) + (\mathbf{6}_c, \mathbf{4}) + (\mathbf{15}_c, \mathbf{2}) \\ \mathbf{120}_{cs} &= (\mathbf{3}_c, \mathbf{2}) + (\mathbf{3}_c, \mathbf{4}) + (\mathbf{15}_c, \mathbf{2}) + (\mathbf{15}_c, \mathbf{4}) + (\mathbf{6}_c, \mathbf{2}). \end{aligned} \quad (74)$$

And the cluster  $q^2\bar{q}$  can be in the flavor representation  $\bar{\mathbf{3}}_f \otimes \bar{\mathbf{3}}_f = \mathbf{3}_f \oplus \bar{\mathbf{6}}_f$  or in  $\mathbf{6}_f \otimes \bar{\mathbf{3}}_f = \mathbf{3}_f \oplus \mathbf{15}_f$ . We can easily see the possible states of the cluster  $q^2\bar{q}$ , which are listed in the followings.

$$\begin{aligned}
& |\mathbf{120}_{cs}, \mathbf{3}_c, \mathbf{2}, \mathbf{3}_f \oplus \bar{\mathbf{6}}_f\rangle, \quad |\mathbf{120}_{cs}, \mathbf{3}_c, \mathbf{4}, \mathbf{3}_f \oplus \bar{\mathbf{6}}_f\rangle, \quad |\mathbf{120}_{cs}, \mathbf{15}_c, \mathbf{2}, \mathbf{3}_f \oplus \bar{\mathbf{6}}_f\rangle, \\
& |\mathbf{120}_{cs}, \mathbf{15}_c, \mathbf{4}, \mathbf{3}_f \oplus \bar{\mathbf{6}}_f\rangle, \quad |\mathbf{120}_{cs}, \bar{\mathbf{6}}_c, \mathbf{2}, \mathbf{3}_f \oplus \bar{\mathbf{6}}_f\rangle, \quad |\mathbf{6}_{cs}, \mathbf{3}_c, \mathbf{2}, \mathbf{3}_f \oplus \bar{\mathbf{6}}_f\rangle, \\
& |\bar{\mathbf{84}}_{cs}, \mathbf{3}_c, \mathbf{2}, \mathbf{3}_f \oplus \mathbf{15}_f\rangle, \quad |\bar{\mathbf{84}}_{cs}, \mathbf{3}_c, \mathbf{4}, \mathbf{3}_f \oplus \mathbf{15}_f\rangle, \quad |\bar{\mathbf{84}}_{cs}, \bar{\mathbf{6}}_c, \mathbf{2}, \mathbf{3}_f \oplus \mathbf{15}_f\rangle \\
& |\bar{\mathbf{84}}_{cs}, \bar{\mathbf{6}}_c, \mathbf{4}, \mathbf{3}_f \oplus \mathbf{15}_f\rangle, \quad |\bar{\mathbf{84}}_{cs}, \mathbf{15}_c, \mathbf{2}, \mathbf{3}_f \oplus \mathbf{15}_f\rangle, \quad |\mathbf{6}_{cs}, \mathbf{3}_c, \mathbf{2}, \mathbf{3}_f \oplus \mathbf{15}_f\rangle.
\end{aligned} \tag{75}$$

Similarly the possible states of another cluster  $q\bar{q}^2$  are

$$\begin{aligned}
& |\bar{\mathbf{120}}_{cs}, \bar{\mathbf{3}}_c, \mathbf{2}, \bar{\mathbf{3}}_f \oplus \mathbf{6}_f\rangle, \quad |\bar{\mathbf{120}}_{cs}, \bar{\mathbf{3}}_c, \mathbf{4}, \bar{\mathbf{3}}_f \oplus \mathbf{6}_f\rangle, \quad |\bar{\mathbf{120}}_{cs}, \bar{\mathbf{15}}_c, \mathbf{2}, \bar{\mathbf{3}}_f \oplus \mathbf{6}_f\rangle, \\
& |\bar{\mathbf{120}}_{cs}, \bar{\mathbf{15}}_c, \mathbf{4}, \bar{\mathbf{3}}_f \oplus \mathbf{6}_f\rangle, \quad |\bar{\mathbf{120}}_{cs}, \mathbf{6}_c, \mathbf{2}, \bar{\mathbf{3}}_f \oplus \mathbf{6}_f\rangle, \quad |\bar{\mathbf{6}}_{cs}, \bar{\mathbf{3}}_c, \mathbf{2}, \bar{\mathbf{3}}_f \oplus \mathbf{6}_f\rangle, \\
& |\mathbf{84}_{cs}, \bar{\mathbf{3}}_c, \mathbf{2}, \bar{\mathbf{3}}_f \oplus \bar{\mathbf{15}}_f\rangle, \quad |\mathbf{84}_{cs}, \bar{\mathbf{3}}_c, \mathbf{4}, \bar{\mathbf{3}}_f \oplus \bar{\mathbf{15}}_f\rangle, \quad |\mathbf{84}_{cs}, \mathbf{6}_c, \mathbf{2}, \bar{\mathbf{3}}_f \oplus \bar{\mathbf{15}}_f\rangle \\
& |\mathbf{84}_{cs}, \mathbf{6}_c, \mathbf{4}, \bar{\mathbf{3}}_f \oplus \bar{\mathbf{15}}_f\rangle, \quad |\mathbf{84}_{cs}, \bar{\mathbf{15}}_c, \mathbf{2}, \bar{\mathbf{3}}_f \oplus \bar{\mathbf{15}}_f\rangle, \quad |\bar{\mathbf{6}}_{cs}, \bar{\mathbf{3}}_c, \mathbf{2}, \bar{\mathbf{3}}_f \oplus \bar{\mathbf{15}}_f\rangle
\end{aligned} \tag{76}$$

We now consider the multiplets whose two subsystems are connected by  $\bar{\mathbf{6}}_c(\mathbf{6}_c)$  or  $\mathbf{15}_c(\bar{\mathbf{15}}_c)$  bond, neglecting these with the clusters being in  $\mathbf{3}_c$  or  $\bar{\mathbf{3}}_c$ , since color triplet bond is easily broken. The states in Eq.(75) and Eq.(76) are eigenstates of the hamiltonian  $H'$ , and the eigenvalues can be calculated straightforwardly. The results are listed in Table IV.

| States                                                                                                                                                                                                           | Eigenvalues of the hamiltonian $H'$ |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------|
| $ \mathbf{120}_{cs}, \mathbf{15}_c, \mathbf{2}, \mathbf{3}_f \oplus \bar{\mathbf{6}}_f\rangle$ ( or $ \bar{\mathbf{120}}_{cs}, \bar{\mathbf{15}}_c, \mathbf{2}, \bar{\mathbf{3}}_f \oplus \mathbf{6}_f\rangle$ ) | $\frac{4}{3}C$                      |
| $ \mathbf{120}_{cs}, \mathbf{15}_c, \mathbf{4}, \mathbf{3}_f \oplus \bar{\mathbf{6}}_f\rangle$ ( or $ \bar{\mathbf{120}}_{cs}, \bar{\mathbf{15}}_c, \mathbf{4}, \bar{\mathbf{3}}_f \oplus \mathbf{6}_f\rangle$ ) | $-\frac{8}{3}C$                     |
| $ \mathbf{120}_{cs}, \bar{\mathbf{6}}_c, \mathbf{2}, \mathbf{3}_f \oplus \bar{\mathbf{6}}_f\rangle$ ( or $ \bar{\mathbf{120}}_{cs}, \mathbf{6}_c, \mathbf{2}, \bar{\mathbf{3}}_f \oplus \mathbf{6}_f\rangle$ )   | $-8C$                               |
| $ \bar{\mathbf{84}}_{cs}, \mathbf{15}_c, \mathbf{2}, \mathbf{3}_f \oplus \mathbf{15}_f\rangle$ ( or $ \mathbf{84}_{cs}, \bar{\mathbf{15}}_c, \mathbf{2}, \bar{\mathbf{3}}_f \oplus \bar{\mathbf{15}}_f\rangle$ ) | $4C$                                |
| $ \bar{\mathbf{84}}_{cs}, \bar{\mathbf{6}}_c, \mathbf{2}, \mathbf{3}_f \oplus \mathbf{15}_f\rangle$ ( or $ \mathbf{84}_{cs}, \mathbf{6}_c, \mathbf{2}, \bar{\mathbf{3}}_f \oplus \bar{\mathbf{15}}_f\rangle$ )   | $\frac{16}{3}C$                     |
| $ \bar{\mathbf{84}}_{cs}, \bar{\mathbf{6}}_c, \mathbf{4}, \mathbf{3}_f \oplus \mathbf{15}_f\rangle$ ( or $ \mathbf{84}_{cs}, \mathbf{6}_c, \mathbf{4}, \bar{\mathbf{3}}_f \oplus \bar{\mathbf{15}}_f\rangle$ )   | $\frac{4}{3}C$                      |

TABLE IV: The allowed states of the cluster  $q^2\bar{q}$  (or another cluster  $q\bar{q}^2$ ) except those states whose color are in representation  $\mathbf{3}_c$  or  $\bar{\mathbf{3}}_c$ , and the corresponding eigenvalues of the colormagnetic interaction hamiltonian  $H'$ .

The states of the total system are formed by combining the states of the cluster  $q^2\bar{q}$  in Eq.(75) and that of another cluster  $q\bar{q}^2$  in Eq.(76), the allowed states of the system and the

corresponding eigenvalues of the hamiltonian  $H'$  are given in the Table V and Table VI. By means of Table V and Table VI, we discuss the enhancements in the  $(q^2\bar{q}) - (q\bar{q}^2)$  cluster configuration in order:

1.  $\Lambda\bar{\Lambda}$ ,  $p\bar{\Lambda}$  and  $p\bar{p}$  enhancements with spin 0 or 1: From the Table V and Table VI, we can see the multiplet state  $|\mathbf{120}_{cs}, \bar{\mathbf{6}}_c, \mathbf{2}, \mathbf{3}_f \oplus \bar{\mathbf{6}}_f\rangle |\mathbf{120}_{cs}, \mathbf{6}_c, \mathbf{2}, \bar{\mathbf{3}}_f \oplus \mathbf{6}_f\rangle$  has the largest colormagnetic energy  $-16C$ , and its flavors are in  $\mathbf{1}(2)_f \oplus \mathbf{8}_f(4) \oplus \mathbf{10}_f \oplus \mathbf{10}_f \oplus \mathbf{27}_f$ . The spin of these states equals 0 or 1, and with the baryon number being zero. Therefore the multiplet state  $|\mathbf{120}_{cs}, \bar{\mathbf{6}}_c, \mathbf{2}, \mathbf{3}_f \oplus \bar{\mathbf{6}}_f\rangle |\mathbf{120}_{cs}, \mathbf{6}_c, \mathbf{2}, \bar{\mathbf{3}}_f \oplus \mathbf{6}_f\rangle$  can decay to (or couple to) baryon octet plus antibaryon octet. Namely,  $(\Lambda\bar{\Lambda}, p\bar{\Lambda} \text{ and } p\bar{p} \text{ enhancements}) \in |\mathbf{120}_{cs}, \bar{\mathbf{6}}_c, \mathbf{2}, \mathbf{3}_f \oplus \bar{\mathbf{6}}_f\rangle |\mathbf{120}_{cs}, \mathbf{6}_c, \mathbf{2}, \bar{\mathbf{3}}_f \oplus \mathbf{6}_f\rangle$ . These baryonium states may mix with the states in the  $(q^3) - (\bar{q}^3)$  cluster configurations illustrated in the previous subsection. Namely, this mixing can be expressed as follows

$$|i\rangle = \alpha_i |(q^3) - (\bar{q}^3)\rangle_i + \beta_i |(q^2\bar{q}) - (q\bar{q}^2)\rangle_i, \quad (77)$$

where  $i = \{\Lambda\bar{\Lambda}, p\bar{\Lambda}, p\bar{p}, \dots\}$ ,  $\alpha_i$  and  $\beta_i$  are constants,  $|i\rangle$  is  $i^{\text{th}}$  physical baryonium state,  $|(q^3) - (\bar{q}^3)\rangle_i$  is  $i^{\text{th}}$  baryonium state in  $(q^3) - (\bar{q}^3)$  cluster configurations, and  $|(q^2\bar{q}) - (q\bar{q}^2)\rangle_i$  is  $i^{\text{th}}$  baryonium state in  $(q^2\bar{q}) - (q\bar{q}^2)$  cluster configurations. We would like to argue that the physical baryonium states may favor the cluster structure with larger mass defect over that with smaller defect. Therefore, comparing the results of this subsection with ones of the previous subsection, we have

$$\text{to } \Lambda\bar{\Lambda} \text{ enhancement : } |\alpha_{i=\Lambda\bar{\Lambda}}| > |\beta_{i=\Lambda\bar{\Lambda}}|, \quad (78)$$

$$\text{to } p\bar{\Lambda} \text{ enhancement : } |\alpha_{i=p\bar{\Lambda}}| \simeq |\beta_{i=p\bar{\Lambda}}|, \quad (79)$$

$$\text{to } p\bar{p} \text{ enhancement : } |\alpha_{i=p\bar{p}}| < |\beta_{i=p\bar{p}}|. \quad (80)$$

In some recent works in literature[9, 11, 12], the  $p\bar{p}$  enhancement has been treated as a bound state of two clusters. Actually, under the approximation of neglecting van der Waal's force between two color singlet baryons, the clusters as elements of the enhancement are objects with color. The potentials used in [9][11][12] should be due to "color chemistry bonds" that are caused by the unfrozen color degree of freedom in the system in the framework of this paper. The Eq.(80) indicates that the color, spin, flavor structures of  $p\bar{p}$  enhancement are mainly of  $|\mathbf{120}_{cs}, \bar{\mathbf{6}}_c, \mathbf{2}, \mathbf{3}_f \oplus \bar{\mathbf{6}}_f\rangle |\mathbf{120}_{cs}, \mathbf{6}_c, \mathbf{2}, \bar{\mathbf{3}}_f \oplus \mathbf{6}_f\rangle$



TABLE VI: The Continuing of Table V.

| States of the total system                                                                                                      | SU <sub>f</sub> (3) structure                                                                                                    | the Spin<br>Structure | the Mass<br>Splitting |
|---------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------|-----------------------|-----------------------|
| $ \overline{84}_{cs}, \overline{6}_c, 4, 3_f \oplus 15_f\rangle  \overline{120}_{cs}, 6_c, 2, \overline{3}_f \oplus 6_f\rangle$ | $1_f \oplus 8_f(4) \oplus 10_f(3) \oplus \overline{10}_f$<br>$\oplus 27_f(2) \oplus \overline{35}_f$                             | <b>1, 2</b>           | $-\frac{20}{3}C$      |
| $ \overline{84}_{cs}, \overline{6}_c, 4, 3_f \oplus 15_f\rangle  84_{cs}, 6_c, 2, \overline{3}_f \oplus \overline{15}_f\rangle$ | $1_f(2) \oplus 8_f(5) \oplus 10_f(2) \oplus \overline{10}(2)$<br>$\oplus 27_f(4) \oplus 35_f \oplus \overline{35}_f \oplus 64_f$ | <b>1, 2</b>           | $\frac{20}{3}C$       |
| $ \overline{84}_{cs}, \overline{6}_c, 4, 3_f \oplus 15_f\rangle  84_{cs}, 6_c, 4, \overline{3}_f \oplus \overline{15}_f\rangle$ | $1_f(2) \oplus 8_f(5) \oplus 10_f(2) \oplus \overline{10}(2)$<br>$\oplus 27_f(4) \oplus 35_f \oplus \overline{35}_f \oplus 64_f$ | <b>0, 1, 2, 3</b>     | $\frac{8}{3}C$        |

in the configuration  $(q^2\overline{q}) - (q\overline{q}^2)$ , and mix with few  $|\overline{70}_{cs}, 8_c, 2, 8_f\rangle |\overline{70}_{cs}, 8_c, 2, 8_f\rangle$  in the configuration  $(q^3) - (\overline{q}^3)$ .

Similarly, Eq.(78) indicates the color-spin-flavor structures of  $\Lambda\overline{\Lambda}$  enhancement are mainly  $|\overline{56}_{cs}, 8_c, 2; 1_f\rangle |\overline{56}_{cs}, 8_c, 2; 1_f\rangle$  in the configuration  $(q^3) - (\overline{q}^3)$ , and plus few mixtures of the corresponding  $\Lambda\overline{\Lambda}$  enhancement state in the configuration  $(q^2\overline{q}) - (q\overline{q}^2)$ . Moreover, by Eq.(79), the  $p\overline{\Lambda}$  enhancement should be half of the states in the  $(q^3) - (\overline{q}^3)$  configuration plus half of that in the  $(q^2\overline{q}) - (q\overline{q}^2)$  approximately.

2. The enhancements with spin -1 or -2: The states  $|\overline{120}_{cs}, \overline{6}_c, 2, 3_f \oplus \overline{6}_f\rangle |84_{cs}, 6_c, 4, \overline{3}_f \oplus \overline{15}_f\rangle$  or  $|\overline{84}_{cs}, \overline{6}_c, 4, 3_f \oplus 15_f\rangle |\overline{120}_{cs}, 6_c, 2, \overline{3}_f \oplus 6_f\rangle$  has colormagnetic energy  $-\frac{20}{3}C$ , but their spins are **1** or **2**. We address that the spins of the baryoniums come from the quark's intrinsic spin in this quark model with  $S$ -wave, instead of coming from their relative orbital angular momentum. These color singlet states are allowed.

On all accounts, both the analysis from common quark model and from the model with quark correlation imply that the  $p\overline{p}$  enhancement,  $p\overline{\Lambda}$  enhancement and the  $\Lambda\overline{\Lambda}$  enhancement are relatively easy to be found and these enhancements which have been seen by BES and Bell are not experiment artifacts. We predict that spin-1  $\Lambda\overline{\Lambda}$  enhancement may be observed in future and we conjecture that BES has observed both spin-0, spin-1  $p\overline{p}$  enhancement and spin-0, spin-1  $p\overline{\Lambda}$  enhancement in section III, these predictions are compatible with the above results obtained in subsection IV.A

and subsection IV.B. These results should be further checked by experiment.

## V. CONCLUSION AND DISCUSSION

In summary, we have studied the mass spectrum of the  $q^3\bar{q}^3$  mesons and the baryon antibaryon enhancements in the quark model with the interaction between quarks being the well-known colormagnetic interaction. First of all, We study  $q^3\bar{q}^3$  system using common quark model without quark correlation with colormagnetic hyperfine interactions between quarks. By this model, we firstly found that the configurations corresponding to the  $\Lambda\bar{\Lambda}$ ,  $p\bar{\Lambda}$ ,  $p\bar{p}$ ,  $\dots$  enhancement can have rather large negative colormagnetic energy. This indicates that these states should be ones with rather low energy, and hence be visible in experiments. These predictions are consistent with the experiments reported in literatures. Secondly, we found further that among those enhancements, the binding colormagnetic energy of spin-0 and spin-1 states are larger than ones of the states with spin-2 and spin-3. Thirdly, the existence of the spin-1  $\Lambda\bar{\Lambda}$ ,  $p\bar{\Lambda}$  and  $p\bar{p}$  enhancement is predicted.

We further study this system in the quark model with quark correlation, we consider two interesting configurations  $(q^3) - (\bar{q}^3)$  and  $(q^2\bar{q}) - (q\bar{q}^2)$ , and the two clusters are connected by color **8**-plet bond, **10**-plet bond and color **6**-plet bond, **15**-plet bond respectively. We find in the spectrum of both configurations there exist rather stable states which has the same quantum number with the  $p\bar{p}$  enhancement,  $p\bar{\Lambda}$  enhancement and  $\Lambda\bar{\Lambda}$  enhancement, hence these enhancements are not experimental artifacts.

To the  $p\bar{p}$  enhancement, we found out that under the approximation of neglecting van der Waal's force between color singlet baryon and color singlet antibaryon, the clusters as elements of the enhancement are objects with color. The potentials used in the studies on the  $p\bar{p}$ -baryonium should be due to "color chemistry bonds" that are caused by the unfrozen color degree of freedom in the system in the framework of this paper. It has been shown that if there exist quark correlation in the baryonium, the state of  $p\bar{p}$  enhancement are mainly of  $|\mathbf{120}_{cs}, \bar{\mathbf{6}}_c, \mathbf{2}, \mathbf{3}_f \oplus \bar{\mathbf{6}}_f\rangle|\mathbf{120}_{cs}, \mathbf{6}_c, \mathbf{2}, \bar{\mathbf{3}}_f \oplus \mathbf{6}_f\rangle$  in the configuration  $(q^2\bar{q}) - (q\bar{q}^2)$ , with small mixing  $|\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}, \mathbf{8}_f\rangle|\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}, \mathbf{8}_f\rangle$  in the configuration  $(q^3) - (\bar{q}^3)$ . Similarly, the state of  $\Lambda\bar{\Lambda}$  enhancement are mainly  $|\mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f\rangle|\mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f\rangle$  in the configuration  $(q^3) - (\bar{q}^3)$ , and plus few mixtures of the corresponding  $\Lambda\bar{\Lambda}$  enhancement state in the configuration  $(q^2\bar{q}) - (q\bar{q}^2)$ . Moreover, the  $p\bar{\Lambda}$  enhancement should be half of the states in the  $(q^3) - (\bar{q}^3)$



configuration plus half of that in the  $(q^2\bar{q}) - (q\bar{q}^2)$  approximately.

Considering all facts from common quark model and from the models with quark correlations, we conclude that the  $p\bar{p}$  enhancement,  $p\bar{\Lambda}$  enhancement and the  $\Lambda\bar{\Lambda}$  enhancement are in existence, and the signal corresponding them must be visible. Actually, they have been shown experimentally by BES and Bell. In other words, those experiments are certainly not experimental artifacts. We predict that spin-**1**  $\Lambda\bar{\Lambda}$  enhancement will be observed in future and we conjecture that BES has observed both spin-**0**, spin-**1**  $p\bar{p}$  enhancement and spin-**0**, spin-**1**  $p\bar{\Lambda}$  enhancement.

The existed quark model with colormagnetic interaction only consider one gluon exchange interaction between quarks(antiquarks) and quarks(antiquarks), and the corrections from the quark's annihilation contribution( the so-called OZI violated graphs ) are not taken into account in the model's framework. The latter will bring small errors to the model, and the problem how to count the contributions of the quark's annihilation diagrams into the model still remains to be open so far. However, since many successes and reasonable results have been achieved in the studies of the multiquark systems within this model framework (i.e., the model is supported by the phenomenology), and our results presented in this paper are exact in the model, we argue that the results and predictions presented in this paper are reliable, and hence should be useful and meaningful for understanding the physics of both  $q^3\bar{q}^3$  system and the recent corresponding experiments reported by BES and Bell qualitatively.

The effect of  $SU(3)_f$  symmetry broken is not taken into account in this paper, and the results are qualitatively correct. It is necessary to consider the effect of strange quark mass in order to make serious comparison with the experiments and predict exactly the masses of the multiplets of the  $q^3\bar{q}^3$  meson.

## ACKNOWLEDGMENTS

This work is partially supported by National Natural Science Foundation of China under Grant Numbers 90403021, and by the PhD Program Funds 20020358040 of the Education Ministry of China and KJCX2-SW-N10 of the Chinese Academy.

---

[1] T.Nakano et al(The LEPS collaboration),Phys.Rev.Lett.91(2003)012002,hep-ex/0301020; BABAR Collaboration, B.Aubert et al., Phys.Rev.Lett.90,242001(2003); Belle Collaboration,

- K.Abe et al., Phys.Rev.Lett.92,012002(2004).
- [2] BES Collaboration, J.Z.Bai et al., Phys.Rev.Lett.91,022001(2003).
  - [3] BES Collaboration, M. Ablikim et al., Phys. Rev. Lett. 95 (2005) 262001; hep-ex/0508025.
  - [4] BES Collaboration, M. Ablikim et al., Phys.Rev. D71 (2005) 072006, hep-ex/0503030.
  - [5] BES Collaboration, M. Ablikim et al., Phys.Rev.Lett.93,112002(2004).
  - [6] Belle Collaboration, K.Abe et al., Phys.Rev.Lett.88,181803(2002).
  - [7] Belle Collaboration, M.-Z.Wang et al., Phys.Rev.Lett.90,201802(2003).
  - [8] Belle Collaboration, Y.-J.Lee et al., Phys.Rev.Lett.93,211801(2004).
  - [9] A.Datta and P.J.O'Donnell, Phys.Lett.B 567,273(2003), hep-ph/0306097.
  - [10] B.S.Zou and H.C.Chiang, Phys.Rev.D69,034004(2004); Xiao-Gang He, Xue-Qian Li and J.P.Ma, Phys.Rev.D71(2005)014031, hep-ph/0407083; Shi-Lin Zhu and Chong-Shou Gao, hep-ph/0507050; C.H.Chang and H.R.Pang, Commun.Theor.Phys.43, 275 (2005), hep-ph/ 0407188; Xiao-Gang He, Xue-Qian Li et al., hep-ph/0509140 ; N.Kochelev and Dong-Pil Min, hep-ph/0508288.
  - [11] Mu-Lin Yan, Si Li, Bin Wu and Bo-Qiang Ma, Phys. Rev. **D72**, 034027 (2005).
  - [12] Gui-Jun Ding and Mu-Lin Yan, Phys.Rev.C72:015208,2005, hep-ph/0502127.
  - [13] J.L.Rosner, Phys.Rev. D68 (2003) 014004, hep-ph/0303079; C.Z.Yuan, X.H.Mo and P.Wang, Phys.Lett.B626 (2005)95-100, hep-ph/0506019.
  - [14] Hong-Mo Chan, Nucl.Phys.B136:401,1978; Phys.Lett.B76:634-640,1978.
  - [15] R.L.Jaffe and F.Wilcek, Phys.Rev.Lett 91,232003(2003), hep-ph/0307341; M.Karliner and H.J.Lipkin, Phys.Lett.B575:249-255,2003, hep-ph/0402260.
  - [16] Gui-Jun Ding and Mu-Lin Yan, Phys.Rev.**D72**, 034014,(2005); hep-ph/0506119.
  - [17] H.Hogaasen and P.Sorba, Nucl.Phys.B145:119,1978; Mod.Phys.Lett.A19:2403-2410,2004, hep-ph/0406078; Fl. Stancu, Phys.Lett. B595 (2004) 269-276; Erratum-ibid. B598 (2004) 295-295, hep-ph/0402044; Carl E. Carlson et al., Phys.Lett. B573 (2003) 101-108, hep-ph/0307396; Jialun Ping et al., Phys.Lett. B602 (2004) 197-204, hep-ph/0408176.
  - [18] A.De Rujula, H.Georgi and S.L.Glashow, Phys,Rev.D.12,147(1975).
  - [19] R.L.Jaffe and K.Johnson, Phys.Lett.B60,201(1976); R.L.Jaffe, Phys.Rev.D15:267,1977; R.L.Jaffe, Phys.Rev.D15:281,1977.
  - [20] P. J. Mulders, A. T. M. Aerts and J. J. De Swart, Phys.Rev.D21, 2653 (1980).
  - [21] D.Strottman, Phys.Rev.D20,748 (1979).

- [22] R.L.Jaffe, hep-ph/0507149.
- [23] Particle Data Group, *14. Quark Model*, in Phys. Lett. **B592**, 1 (2004), *pp154-159*.
- [24] C.Itzykson, M.Nauenberg, Rev.Mod.Phys.38:95-120,1966.
- [25] S. I. So and D. Strottman, J.Math.Phys.20:153-176, 1979; D. Strottman, J.Math.Phys.20:1643-1647,1979.
- [26] Jin-Quan Chen et al, Tables of the  $SU(mn) \supset SU(m) \times SU(n)$  Coefficients of Fractional Parentage (World Scientific, 1991).
- [27] W.G.McKay and J.Patera, Tables of Dimensions, Indices, and Branching Rules for Representations of Simple Lie Algebras, (Marcel Dekker, New York, 1981) p.98.

## APPENDIX A: THE MIXING OF STATES WITH DEFINITE SPIN UNDER THE COLORMAGNETIC INTERACTION HAMILTONIAN $H'$

In this appendix, as an example, we will illustrate that in the case of  $70_{cs} \otimes \overline{20}_{cs}$  the three spin-1 states  $|35_{cs}, 1_c, 3; 8_f \otimes \overline{10}_f\rangle$ ,  $|280_{cs}, 1_c, 3; 8_f \otimes \overline{10}_f\rangle$  and  $|896_{cs}, 1_c, 3; 8_f \otimes \overline{10}_f\rangle$  are mixed due to the colormagnetic interaction  $H'$ , and are linear combinations of the states  $|(\overline{70}_{cs}, 8_c, 4; 8_f), (\overline{20}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 3, 8_f \otimes \overline{10}_f\rangle$ ,  $|(\overline{70}_{cs}, 8_c, 2; 8_f)(\overline{20}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 3, 8_f \otimes \overline{10}_f\rangle$  and  $|(\overline{70}_{cs}, 1_c, 2; 8_f), (\overline{20}_{cs}, 1_c, 4; \overline{10}_f), 1_c, 3, 8_f \otimes \overline{10}_f\rangle$ . The eigenstates and the corresponding colormagnetic interaction energy are calculated, for the other cases, these quantities can also be calculated following the same method.

$$\begin{aligned}
|896_{cs}, 1_c, 3; 8_f \otimes \overline{10}_f\rangle &= \left\{ \begin{array}{cc|c} 70_{cs} & \overline{20}_{cs} & 896_{cs} \\ (8_c, 4) & (8_c, 2) & (1_c, 3) \end{array} \right\} |(\overline{70}_{cs}, 8_c, 4; 8_f), (\overline{20}_{cs}, 8_c, 2; \overline{10}_f), \\
&1_c, 3; 8_f \otimes \overline{10}_f\rangle + \left\{ \begin{array}{cc|c} 70_{cs} & \overline{20}_{cs} & 896_{cs} \\ (8_c, 2) & (8_c, 2) & (1_c, 3) \end{array} \right\} |(\overline{70}_{cs}, 8_c, 2; 8_f), (\overline{20}_{cs}, 8_c, 2; \overline{10}_f), \\
&1_c, 3; 8_f \otimes \overline{10}_f\rangle + \left\{ \begin{array}{cc|c} 70_{cs} & \overline{20}_{cs} & 896_{cs} \\ (1_c, 2) & (1_c, 4) & (1_c, 3) \end{array} \right\} |(\overline{70}_{cs}, 1_c, 2; 8_f), (\overline{20}_{cs}, 1_c, 4; \overline{10}_f), \\
&1_c, 3; 8_f \otimes \overline{10}_f\rangle; \\
|280_{cs}, 1_c, 3; 8_f \otimes \overline{10}_f\rangle &= \left\{ \begin{array}{cc|c} 70_{cs} & \overline{20}_{cs} & 280_{cs} \\ (8_c, 4) & (8_c, 2) & (1_c, 3) \end{array} \right\} |(\overline{70}_{cs}, 8_c, 4; 8_f), (\overline{20}_{cs}, 8_c, 2; \overline{10}_f),
\end{aligned}$$

$$\begin{aligned}
& \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f \rangle + \left\{ \begin{array}{cc|c} \mathbf{70}_{cs} & \overline{\mathbf{20}}_{cs} & \mathbf{280}_{cs} \\ (8_c, 2) & (8_c, 2) & (1_c, 3) \end{array} \right\} |(\mathbf{70}_{cs}, 8_c, 2; \mathbf{8}_f), (\overline{\mathbf{20}}_{cs}, 8_c, 2; \overline{\mathbf{10}}_f), \\
& \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f \rangle + \left\{ \begin{array}{cc|c} \mathbf{70}_{cs} & \overline{\mathbf{20}}_{cs} & \mathbf{280}_{cs} \\ (1_c, 2) & (1_c, 4) & (1_c, 3) \end{array} \right\} |(\mathbf{70}_{cs}, 1_c, 2; \mathbf{8}_f), (\overline{\mathbf{20}}_{cs}, 1_c, 4; \overline{\mathbf{10}}_f), \\
& \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f \rangle; \\
& |\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f \rangle = \left\{ \begin{array}{cc|c} \mathbf{70}_{cs} & \overline{\mathbf{20}}_{cs} & \mathbf{35}_{cs} \\ (8_c, 4) & (8_c, 2) & (1_c, 3) \end{array} \right\} |(\mathbf{70}_{cs}, 8_c, 4; \mathbf{8}_f), (\overline{\mathbf{20}}_{cs}, 8_c, 2; \overline{\mathbf{10}}_f), \\
& \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f \rangle + \left\{ \begin{array}{cc|c} \mathbf{70}_{cs} & \overline{\mathbf{20}}_{cs} & \mathbf{35}_{cs} \\ (8_c, 2) & (8_c, 2) & (1_c, 3) \end{array} \right\} |(\mathbf{70}_{cs}, 8_c, 2; \mathbf{8}_f), (\overline{\mathbf{20}}_{cs}, 8_c, 2; \overline{\mathbf{10}}_f), \\
& \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f \rangle + \left\{ \begin{array}{cc|c} \mathbf{70}_{cs} & \overline{\mathbf{20}}_{cs} & \mathbf{35}_{cs} \\ (1_c, 2) & (1_c, 4) & (1_c, 3) \end{array} \right\} |(\mathbf{70}_{cs}, 1_c, 2; \mathbf{8}_f), (\overline{\mathbf{20}}_{cs}, 1_c, 4; \overline{\mathbf{10}}_f), \\
& \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f \rangle; \tag{A1}
\end{aligned}$$

$\text{SU}_{cs}(6) \supset \text{SU}_c(3) \times \text{SU}_s(2)$  coefficients of fractional parentage can be calculated following the Ref.[25], and some results has been listed in the Ref. [26]. From this book, we can find the  $\text{SU}_{cs}(6) \supset \text{SU}_c(3) \times \text{SU}_s(2)$  coefficients of fractional parentage which appear in the above formula,

$$\begin{aligned}
& \left\{ \begin{array}{cc|c} \mathbf{70}_{cs} & \overline{\mathbf{20}}_{cs} & \mathbf{896}_{cs} \\ (8_c, 4) & (8_c, 2) & (1_c, 3) \end{array} \right\} = -\sqrt{\frac{1}{9}}, \quad \left\{ \begin{array}{cc|c} \mathbf{70}_{cs} & \overline{\mathbf{20}}_{cs} & \mathbf{896}_{cs} \\ (8_c, 2) & (8_c, 2) & (1_c, 3) \end{array} \right\} = -\sqrt{\frac{4}{9}} \\
& \left\{ \begin{array}{cc|c} \mathbf{70}_{cs} & \overline{\mathbf{20}}_{cs} & \mathbf{896}_{cs} \\ (1_c, 2) & (1_c, 4) & (1_c, 3) \end{array} \right\} = \sqrt{\frac{4}{9}}, \quad \left\{ \begin{array}{cc|c} \mathbf{70}_{cs} & \overline{\mathbf{20}}_{cs} & \mathbf{280}_{cs} \\ (8_c, 4) & (8_c, 2) & (1_c, 3) \end{array} \right\} = -\sqrt{\frac{4}{9}} \\
& \left\{ \begin{array}{cc|c} \mathbf{70}_{cs} & \overline{\mathbf{20}}_{cs} & \mathbf{280}_{cs} \\ (8_c, 2) & (8_c, 2) & (1_c, 3) \end{array} \right\} = -\sqrt{\frac{1}{9}}, \quad \left\{ \begin{array}{cc|c} \mathbf{70}_{cs} & \overline{\mathbf{20}}_{cs} & \mathbf{280}_{cs} \\ (1_c, 2) & (1_c, 4) & (1_c, 3) \end{array} \right\} = -\sqrt{\frac{4}{9}} \\
& \left\{ \begin{array}{cc|c} \mathbf{70}_{cs} & \overline{\mathbf{20}}_{cs} & \mathbf{35}_{cs} \\ (8_c, 4) & (8_c, 2) & (1_c, 3) \end{array} \right\} = -\sqrt{\frac{4}{9}}, \quad \left\{ \begin{array}{cc|c} \mathbf{70}_{cs} & \overline{\mathbf{20}}_{cs} & \mathbf{35}_{cs} \\ (8_c, 2) & (8_c, 2) & (1_c, 3) \end{array} \right\} = \sqrt{\frac{4}{9}} \\
& \left\{ \begin{array}{cc|c} \mathbf{70}_{cs} & \overline{\mathbf{20}}_{cs} & \mathbf{35}_{cs} \\ (1_c, 2) & (1_c, 4) & (1_c, 3) \end{array} \right\} = \sqrt{\frac{1}{9}} \tag{A2}
\end{aligned}$$

Substituting Eq.(A2) into Eq.(A1) we can obtain

$$|\mathbf{896}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f \rangle = -\sqrt{\frac{1}{9}} |(\mathbf{70}_{cs}, 8_c, 4; \mathbf{8}_f), (\overline{\mathbf{20}}_{cs}, 8_c, 2; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f \rangle$$

$$\begin{aligned}
& -\sqrt{\frac{4}{9}}|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle + \sqrt{\frac{4}{9}}|(\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), \\
& (\overline{\mathbf{20}}_{cs}, \mathbf{1}_c, \mathbf{4}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle; \\
& |\mathbf{280}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle = -\sqrt{\frac{4}{9}}|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle \\
& -\sqrt{\frac{1}{9}}|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle - \sqrt{\frac{4}{9}}|(\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), \\
& (\overline{\mathbf{20}}_{cs}, \mathbf{1}_c, \mathbf{4}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle; \\
& |\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle = -\sqrt{\frac{4}{9}}|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle \\
& +\sqrt{\frac{4}{9}}|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle + \sqrt{\frac{1}{9}}|(\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), \\
& (\overline{\mathbf{20}}_{cs}, \mathbf{1}_c, \mathbf{4}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle
\end{aligned} \tag{A3}$$

From the above, we can learn that  $\langle \mathbf{896}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f | H' | \mathbf{896}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_{cs} \rangle = \frac{516}{27}C$ ,  $\langle \mathbf{896}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f | H' | \mathbf{280}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_{cs} \rangle = \frac{90}{9}C$  etc., so  $H'$  can be expressed as the the following matrix form in the space expanded by the three spin-1 states  $|\mathbf{896}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8} \otimes \overline{\mathbf{10}}_f\rangle$ ,  $|\mathbf{280}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8} \otimes \overline{\mathbf{10}}_f\rangle$ ,  $|\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8} \otimes \overline{\mathbf{10}}_f\rangle$

$$\frac{1}{27}C \begin{pmatrix} 516 & 240 & -48 \\ 240 & 264 & 192 \\ -48 & 192 & -240 \end{pmatrix} \tag{A4}$$

Diagonalizing the above matrix, we can obtain the eigenstates and the corresponding color-magnetic interaction energy,

$$\begin{aligned}
& |\mathbf{3}, \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle_1 = 0.841|\mathbf{896}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle + 0.536|\mathbf{280}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle \\
& +0.069|\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle = -0.684|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle \\
& -0.693|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle + 0.226|(\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), \\
& (\overline{\mathbf{20}}_{cs}, \mathbf{1}_c, \mathbf{4}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle,
\end{aligned} \tag{A5}$$

$$\begin{aligned}
& |\mathbf{3}, \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle_2 = 0.156|\mathbf{896}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle - 0.364|\mathbf{280}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle \\
& +0.918|\mathbf{35}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle = -0.422|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle \\
& +0.629|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{20}}_{cs}, \mathbf{8}_c, \mathbf{2}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle + 0.653|(\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), \\
& (\overline{\mathbf{20}}_{cs}, \mathbf{1}_c, \mathbf{4}; \overline{\mathbf{10}}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle,
\end{aligned} \tag{A6}$$

$$|\mathbf{3}, \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle_3 = -0.518|\mathbf{896}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle + 0.762|\mathbf{280}_{cs}, \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \overline{\mathbf{10}}_f\rangle$$

$$\begin{aligned}
& +0.389|35_{cs}, 1_c, 3; 8_f \otimes \overline{10}_f\rangle = -0.595|(\overline{70}_{cs}, 8_c, 4; 8_f), (\overline{20}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle \\
& +0.351|(\overline{70}_{cs}, 8_c, 2; 8_f), (\overline{20}_{cs}, 8_c, 2; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle - 0.723|(\overline{70}_{cs}, 1_c, 2; 8_f), \\
& (\overline{20}_{cs}, 1_c, 4; \overline{10}_f), 1_c, 3; 8_f \otimes \overline{10}_f\rangle
\end{aligned} \tag{A7}$$

The colormagnetic energy respectively are 24.634C, -12.007C, 7.373C.

## APPENDIX B: THE MATRIX REPRESENTATION OF THE HAMILTONIAN $H'$ IN THE SUBSPACE OF STATES WITH FIXED SPIN

In the following, we list the matrix form of the colormagnetic interaction hamiltonian  $H'$  in the subspace expanded by the states of fixed spin in each case, and the constant  $C$  in front each of matrix is omitted.

### 1. $56_{cs} \otimes \overline{56}_{cs}$

A), the spin-0 state, there exist two spin-0 states  $|(\overline{56}_{cs}, 10_c, 4; 1_f), (\overline{56}_{cs}, \overline{10}_c, 4; 1_f), 1_c, 1; 1_f \otimes 1_f\rangle$  and  $|(\overline{56}_{cs}, 8_c, 2; 1_f), (\overline{56}_{cs}, 8_c, 2; 1_f), 1_c, 1; 1_f \otimes 1_f\rangle$ , and the hamiltonian  $H'$  can be expressed as the following matrix form in the subspace expanded by the latter state vectors ,

$$\begin{pmatrix} -48 & -8\sqrt{10} \\ -8\sqrt{10} & -64 \end{pmatrix} \tag{B1}$$

B), the spin-1 state, in the subspace expanded by the two spin-1 states  $|(\overline{56}_{cs}, 10_c, 4; 1_f), (\overline{56}_{cs}, \overline{10}_c, 4; 1_f), 1_c, 3; 1_f \otimes 1_f\rangle$  and  $|(\overline{56}_{cs}, 8_c, 2; 1_f), (\overline{56}_{cs}, 8_c, 2; 1_f), 1_c, 3; 1_f \otimes 1_f\rangle$ , the matrix representation of  $H'$  is

$$\begin{pmatrix} -112/3 & -40\sqrt{2}/3 \\ -40\sqrt{2}/3 & -16 \end{pmatrix} \tag{B2}$$

### 2. $70_{cs} \otimes \overline{56}_{cs}$ and $56_{cs} \otimes \overline{70}_{cs}$ .

A), the spin-1 state, the three spin-1 states  $|(\overline{70}_{cs}, 10_c, 2; 8_f), (\overline{56}_{cs}, \overline{10}_c, 4; 1_f), 1_c, 3; 8_f \otimes 1_f\rangle$ ,  $|(\overline{70}_{cs}, 8_c, 4; 8_f), (\overline{56}_{cs}, 8_c, 2; 1_f), 1_c, 3; 8_f \otimes 1_f\rangle$ , and  $|(\overline{70}_{cs}, 8_c, 2; 8_f), (\overline{56}_{cs}, 8_c, 2; 1_f), 1_c, 3; 8_f \otimes 1_f\rangle$  expand the Hilbert space,

and the matrix form of  $H'$  is

$$\begin{pmatrix} -40/3 & 4\sqrt{10}/3 & 16\sqrt{10}/3 \\ 4\sqrt{10}/3 & -76/3 & -56/3 \\ 16\sqrt{10}/3 & -56/3 & -40/3 \end{pmatrix} \quad (\text{B3})$$

B), the spin-2 state, there are two spin-2 base vectors  $|(\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; \mathbf{8}_f), (\mathbf{56}_{cs}, \mathbf{10}_c, \mathbf{4}; \mathbf{1}_f), \mathbf{1}_c, \mathbf{5}; \mathbf{8}_f \otimes \mathbf{1}_f\rangle$  and  $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\mathbf{56}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{1}_f), \mathbf{1}_c, \mathbf{5}; \mathbf{8}_f \otimes \mathbf{1}_f\rangle$ , and  $H'$ 's matrix form is the following form:

$$\begin{pmatrix} 8 & 4\sqrt{10} \\ 4\sqrt{10} & -4 \end{pmatrix} \quad (\text{B4})$$

### 3. $\mathbf{70}_{cs} \otimes \mathbf{70}_{cs}$

A), the spin-2 states, there are three spin-2 states  $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{5}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ ,  $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{5}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle$  and  $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{5}, \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ , and the hamiltonian  $H'$  can be expressed as the following matrix form in the subspace expanded by them,

$$\begin{pmatrix} 0 & -4\sqrt{2} & -4\sqrt{2} \\ -4\sqrt{2} & 4 & -12 \\ -4\sqrt{2} & -12 & 4 \end{pmatrix} \quad (\text{B5})$$

B), the spin-1 states, in the subspace of six spin-1 states  $|(\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; \mathbf{8}_f), (\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ ,  $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ ,  $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ ,  $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ ,  $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$  and  $|(\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), (\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{3}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ , the matrix representation of  $H'$  is

$$\begin{pmatrix} \frac{32}{3} & -\frac{20\sqrt{2}}{3} & \frac{16\sqrt{5}}{3} & \frac{16\sqrt{5}}{3} & -\frac{8\sqrt{5}}{3} & 0 \\ -\frac{20\sqrt{2}}{3} & -\frac{32}{3} & -\frac{4\sqrt{10}}{3} & -\frac{4\sqrt{10}}{3} & 4\sqrt{10} & \frac{8\sqrt{5}}{3} \\ \frac{16\sqrt{5}}{3} & -\frac{4\sqrt{10}}{3} & -\frac{20}{3} & -4 & -\frac{8}{3} & \frac{16\sqrt{2}}{3} \\ \frac{16\sqrt{5}}{3} & -\frac{4\sqrt{10}}{3} & -4 & -\frac{20}{3} & -\frac{8}{3} & \frac{16\sqrt{2}}{3} \\ -\frac{8\sqrt{5}}{3} & 4\sqrt{10} & -\frac{8}{3} & -\frac{8}{3} & -\frac{8}{3} & \frac{8\sqrt{2}}{3} \\ 0 & \frac{8\sqrt{5}}{3} & \frac{16\sqrt{2}}{3} & \frac{16\sqrt{2}}{3} & \frac{8\sqrt{2}}{3} & -16 \end{pmatrix} \quad (\text{B6})$$


C), the spin-0 state, the four spin-0 states  $|(\mathbf{70}_{cs}, \mathbf{10}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \overline{\mathbf{10}}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ ,  
 $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{4}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$ ,  
 $|(\mathbf{70}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{8}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$  and  $|(\mathbf{70}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), (\overline{\mathbf{70}}_{cs}, \mathbf{1}_c, \mathbf{2}; \mathbf{8}_f), \mathbf{1}_c, \mathbf{1}; \mathbf{8}_f \otimes \mathbf{8}_f\rangle$  expand the Hilbert space. And the hamiltonian  $H'$  can be expressed as




$$\begin{pmatrix} 0 & -4\sqrt{10} & 8\sqrt{5} & 0 \\ -4\sqrt{10} & -16 & 12\sqrt{2} & 8 \\ 8\sqrt{5} & 12\sqrt{2} & -8 & -8\sqrt{2} \\ 0 & 8 & -8\sqrt{2} & -16 \end{pmatrix} \quad (\text{B7})$$








TABLE VIII: The Continuing of Table VII.




| Young tabular                                                                      | Dimension | $SU_c(3) \otimes SU_s(2)$ content                                                                                                                                                                                                                                                  | Eigenvalue of<br>Casimir operator |
|------------------------------------------------------------------------------------|-----------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------|
|   | 280       | $(\mathbf{1}_c, \mathbf{3}), (\mathbf{8}_c, \mathbf{1}), 2(\mathbf{8}_c, \mathbf{3}), (\mathbf{8}_c, \mathbf{5}), (\mathbf{10}_c, \mathbf{1}), (\mathbf{10}_c, \mathbf{3}), (\mathbf{10}_c, \mathbf{5}),$<br>$(\overline{\mathbf{10}}_c, \mathbf{1}), (\mathbf{27}_c, \mathbf{3})$ | 24                                |
|   | 189       | $(\mathbf{1}_c, \mathbf{1}), (\mathbf{1}_c, \mathbf{5}), (\mathbf{8}_c, \mathbf{1}), 2(\mathbf{8}_c, \mathbf{3}), (\mathbf{8}_c, \mathbf{5}), (\mathbf{10}_c, \mathbf{3}), (\overline{\mathbf{10}}_c, \mathbf{3}),$<br>$(\mathbf{27}_c, \mathbf{1})$                               | 20                                |
|   | 175       | $(\mathbf{1}_c, \mathbf{3}), (\mathbf{1}_c, \mathbf{7}), (\mathbf{8}_c, \mathbf{3}), (\mathbf{8}_c, \mathbf{5}), (\mathbf{10}_c, \mathbf{1}), (\overline{\mathbf{10}}_c, \mathbf{1}), (\mathbf{27}_c, \mathbf{3})$                                                                 | 24                                |
|  | 35        | $(\mathbf{1}_c, \mathbf{3}), (\mathbf{8}_c, \mathbf{1}), (\mathbf{8}_c, \mathbf{3})$                                                                                                                                                                                               | 12                                |

(I)   $\otimes$   =   $\oplus$    $\oplus$    $\oplus$    $\oplus$    $\oplus$    $\oplus$    $\oplus$  

(II)   $\otimes$   =   $\oplus$    $\oplus$    $\oplus$  




(III)   $\otimes$   =   $\oplus$    $\oplus$    $\oplus$  

(IV)   $\otimes$   =   $\oplus$  

(V)   $\otimes$   =   $\oplus$    $\oplus$    $\oplus$  

(VI)   $\otimes$   =   $\oplus$    $\oplus$    $\oplus$  

FIG. 1: The  $SU(6)_{cs}$  outer product for  $q^3$  and  $\bar{q}^3$  (I)  $70_{cs} \otimes \overline{70}_{cs} = 3675_{cs} \oplus 405_{cs} \oplus 280_{cs} \oplus \overline{280}_{cs} \oplus 35_{cs} \oplus 189_{cs} \oplus 35_{cs} \oplus 1_{cs}$ ; (II)  $70_{cs} \otimes \overline{56}_{cs} = 3200_{cs} \oplus 405_{cs} \oplus \overline{280}_{cs} \oplus 35_{cs}$ ; (III)  $70_{cs} \otimes \overline{20}_{cs} = 896_{cs} \oplus 280_{cs} \oplus 189_{cs} \oplus 35_{cs}$ ; (IV)  $56_{cs} \otimes \overline{20}_{cs} = 840_{cs} \oplus 280_{cs}$ ; (V)  $56_{cs} \otimes \overline{56}_{cs} = 2695_{cs} \oplus 405_{cs} \oplus 35_{cs} \oplus 1_{cs}$ ; (VI)  $20_{cs} \otimes \overline{20}_{cs} = 175_{cs} \oplus 189_{cs} \oplus 35_{cs} \oplus 1_{cs}$ .

(I)   $\otimes$   =   $\oplus$    $\oplus$    $\oplus$  



(II)   $\otimes$   =   $\oplus$    $\oplus$    $\oplus$  

FIG. 2: The  $SU(3)_f$  direct product for  $q^3$  and  $\bar{q}^3$  (I)  $8_f \otimes \overline{10}_f = \overline{35}_f \oplus 27_f \oplus 8_f \oplus \overline{10}_f$ ; (II)  $10_f \otimes \overline{10}_f = 64_f \oplus 27_f \oplus 8_f \oplus 1_f$




FIG. 3: The configuration in common quark model, where gray circles denote quarks and red circles denote antiquarks




FIG. 4: The configuration  $(q^3) - (\bar{q}^3)$ , where gray circles denote quarks and red circles denote antiquarks.




FIG. 5: The configuration  $(q^2\bar{q}) - (q\bar{q}^2)$ , where gray circles denote quarks and red circles denote antiquarks